

# The annual number of breeding adults and the effective population size of syntopic newts (*Triturus cristatus*, *T. marmoratus*)

R. JEHLLE,\* J. W. ARNTZEN,† T. BURKE,‡ A. P. KRUPA‡ and W. HÖDL\*

\*Institute of Zoology, University of Vienna, Althanstrasse 14, A-1090 Vienna, Austria, †Universidade do Porto, Unidade de Génética Animal e Conservação, Campus Agrário de Vairão, Rua de Monte, Crasto, 4480 Vila do Conde, Portugal, ‡Department of Animal and Plant Sciences, University of Sheffield, Western Bank, S10 2TN Sheffield, UK

## Abstract

Pond-breeding amphibians are deme-structured organisms with a population genetic structure particularly susceptible to demographic threats. We estimated the effective number of breeding adults ( $N_b$ ) and the effective population size ( $N_e$ ) of the European urodele amphibians *Triturus cristatus* (the crested newt) and *T. marmoratus* (the marbled newt), using temporal shifts in microsatellite allele frequencies. Eight microsatellite loci isolated from a *T. cristatus* library were used, five of which proved polymorphic in *T. marmoratus*, albeit with high frequencies of null alleles at two loci. Three ponds in western France were sampled, situated 4–10 kilometres apart and inhabited by both species. Parent–offspring cohort comparisons were used to measure  $N_b$ ; samples collected at time intervals of nine or 12 years, respectively, were used to measure  $N_e$ . The adult population census size ( $N$ ) was determined by mark–recapture techniques. With one exception, genetic distances ( $F_{ST}$ ) between temporal samples were lower than among populations.  $N_b$  ranged between 10.6 and 101.8 individuals,  $N_e$  ranged between 9.6 and 13.4 individuals. For the pond where both parameters were available,  $N_b/N$  (overall range: 0.10–0.19) was marginally larger than  $N_e/N$  (overall range: 0.09–0.16), which is reflected in the temporal stability of  $N$ . In line with the observed differences in reproductive life-histories between the species,  $N_b/N$  ratios for newts were about one order of magnitude higher than for the anuran amphibian *Bufo bufo*. Despite of the colonization of the study area by *T. cristatus* only some decades ago, no significant genetic bottleneck could be detected. Our findings give rise to concerns about the long-term demographic viability of amphibian populations in situations typical for European landscapes.

**Keywords:** effective number of breeders, effective population size, microsatellites, temporal method, *Triturus cristatus*, *Triturus marmoratus*

Received 21 July 2000; revision received 25 October 2000; accepted 25 October 2000

## Introduction

The concept of effective population size ( $N_e$ ) is at the heart of population genetics.  $N_e$  is the most important determinant for the amount of genetic drift, a fundamental parameter influencing genetic population structure. For conservation biological issues,  $N_e$  is regarded as important because it is the main reason behind loss of genetic diversity (Frankham 1995a), with low  $N_e$  increasing the

probability for population extinction (Newman & Pilson 1997). However, measuring  $N_e$  is notoriously difficult, and not all analytical methods lead to comparable results. Of particular practical importance in the study of free-living organisms is the ratio of  $N_e$  to the adult population census size ( $N$ ). Demographic expectations of  $N_e/N$  range from 0.25 to 0.75, assuming a stable  $N$  (Nunney 1993), whereas empirical studies revealed on average much lower values (0.1, Frankham 1995b).

Genetic estimates of  $N_e$  in wild populations are often obtained with the ‘temporal method’, which is based on temporal variance in marker allele frequencies (Waples

Correspondence: Robert Jehle. Fax: +43 142779544; E-mail: Robert.Jehle@univie.ac.at

1989a; Schwartz *et al.* 1998; Williamson & Slatkin 1999; see also Crandall *et al.* 1999). Microsatellites are currently having a large impact in many studies of evolutionary and conservation biology, as they have enabled genetic variability to be measured and compared with a higher resolution than other markers such as allozymes. However, when measuring the degree of differentiation between populations, the high mutation rate and distinct mutational behaviour of microsatellites requires a careful interpretation of the observed variation (Hedrick 1999), and/or new analytical tools (Luikart & England 1999). In sampling regimes typical for the temporal method the probability impact of mutation events can be regarded as negligible. Nevertheless, few studies have so far measured  $N_e$  from microsatellite genetic variation (but see Miller & Kapuscinski 1997; Lehmann *et al.* 1998; Fiumera *et al.* 2000).

Many temperate amphibians reproduce in small breeding ponds and are particularly amenable to population studies.  $N_e$ , or the related parameter  $N_b$  (the effective number of breeding adults), has been estimated as mostly below 100 individuals (Merrell 1968; Gill 1978; Eastal 1985; Berven & Grudzien 1990; Scribner *et al.* 1997; Driscoll 1999; Funk *et al.* 1999; Seppä & Laurila 1999), but the underlying data sets and the biological significance of the results are based on different methods and, therefore, not easy to interpret.  $N_e$  and  $N_b$  are expected to be low when variance in female reproductive success is high, multiple matings are rare, and the sex ratio is skewed (Nunney 1993; Nunney 1996). Scribner *et al.* (1997) used temporal shifts at DNA (minisatellite) markers between adults and their offspring for estimating the effective number of breeding anurans *Bufo bufo* (common toad), a species characterized by 'explosive' breeding at uneven sex ratio. In fact, they revealed  $N_b/N$  ratios as low as around 0.01.

European urodeles of the genus *Triturus* usually have a sex ratio close to 1:1, with females producing fewer eggs than *B. bufo* and receiving spermatophores from several males (Halliday 1998). In western France, the two large-bodied newts *Triturus cristatus* (crested newt) and *T. marmoratus* (marbled newt) are largely separated by their occurrence in different landscapes, and locally form a hybrid zone where they breed in identical ponds. Due to a recent expansion of the distribution range, *T. cristatus* currently inhabits areas which were some decades ago exclusively occupied by *T. marmoratus* (Arntzen & Wallis 1991). The objectives of this study are to: (i) compare the degree of spatial and temporal genetic differentiation of syntopic *T. cristatus* and *T. marmoratus* populations; (ii) measure  $N_b$  and  $N_b/N$  in both species by comparing microsatellite allele frequencies of adults and their progeny; (iii) measure  $N_e$  and  $N_e/N$  with samples collected at time intervals of nine and 12 years, respectively; (iv) test the hypothesis that newts have a higher  $N_b/N$  ratio than common toads; and (v) test the hypothesis that *T. cristatus* populations

experienced a genetic bottleneck during the recent colonization of the new ponds.

## Materials and methods

### Sample collections

Samples were collected at three cattle ponds inhabited by both study species, situated in the 'Département de Mayenne'. For detailed descriptions, locations, and schematic drawings of Ponds 1 and 3 see Jehle *et al.* (2000). Pond 2 (about 30 m<sup>2</sup> in area and 1 m deep) is identical with Pond 278 in Arntzen & Wallis (1991). The straight-line distance is 3.8 km between Ponds 1 and 2, 10.5 km between Ponds 1 and 3, and 7.3 km between Ponds 2 and 3.

Newts were captured from all pond regions by dipnetting or underwater funnel traps (Jehle *et al.* 2000). Individuals were marked by removing toes. Estimates of  $N \pm SE$  were obtained using mark-recapture calculations (Begon's weighted mean, Begon 1979), which are based on comparisons between numbers of recaptured and newly encountered individuals in successive sampling sessions. In 1998, the toes were stored in 96% ethanol and used as tissue samples for microsatellite analysis. Other samples were obtained from sacrificed larvae collected in 1986 (Pond 2) and 1989 (Pond 1), and nondestructively by removing a gill from larvae captured in 1998 (Ponds 1 and 3).

### Microsatellite genotyping and scoring

DNA extractions were performed using standard phenol-chloroform procedures (Sambrook *et al.* 1989). Primers were used for eight microsatellite loci designed from a *Triturus cristatus* library (Krupa *et al.* in preparation). Polymerase chain reaction (PCR) amplifications were carried out in 10 µL reaction volumes under the following conditions: 1 or 1.5 mM MgCl<sub>2</sub>, 0.1 mM dNTPs, 0.5 U polymerase (Thermoprime Plus, Advanced Biotechnologies, Epsom, UK) and 0.1 mM of each primer in the manufacturer's buffer (buffer IV). Thermal profiles for *Tcri27*, *Tcri35*, and *Tcri43* were 39 cycles of 30 s at 93 °C, 30 s at the primer-specific annealing temperature, 45 s at 72 °C; for *Tcri13*, *Tcri29*, *Tcri32* and *Tcri46b* we used a 'touchdown' thermal profile (for more details see Krupa *et al.* in preparation). *Tcri46b* amplifies the same microsatellite as *Tcri46* in Krupa *et al.* (in preparation), but with a different primer sequence (forward: GTTTGGGTAGCCATGCACTT, reverse: ATCCAAGCATTGGGATTCA). Primers were labelled with fluorochromes TET, HEX, and FAM, and PCR products were run on 5% denaturing polyacrylamide gels (Long Ranger) on an automated ABI 377 DNA sequencer. PCR products were compiled and analysed with ABI GENESCAN software using standard Tamra 2500 (*Tcri32*) or

Tamra 500 (all other primers). Samples for which more than two primers failed to amplify were not included in the analysis.

For scoring the PCR products we used ABI GENOTYPER and visual inspections of gels. Allele sizes were rounded to the nearest integer. Called and actual allele size can differ by about one base pair (Haberl & Tautz 1999), but the tetranucleotide nature of our microsatellites allowed unambiguous designation within and between gels.

#### Data analysis

Observed ( $H_O$ ) and expected ( $H_E$ ) heterozygosity, probability tests for genotypic linkage disequilibrium and measures of genetic differentiation ( $F_{ST}$ ) were computed using GENEPOP 3.1d (Raymond & Rousset 1995). Temporal genetic differences within populations should not be described with conventional contingency tests (Waples 1989b), and we used  $F_{ST}$  for direct comparisons between temporal and spatial degrees of differentiation (cf. Nielsen *et al.* 1999). Departures from Hardy–Weinberg equilibrium at each locus were also computed using GENEPOP, with the Markov chain method (Guo & Thompson 1992) to obtain unbiased estimates of Fisher's exact test through 1000 iterations. Pairwise Wilcoxon tests were used to compare observed heterozygosities and numbers of alleles per locus between temporal samples. Frequencies of null alleles were estimated using the algorithm provided in CERVUS 1.0 (Marshall *et al.* 1998), with data pooled across populations. Tests for recent bottlenecks were conducted with BOTTLENECK 1.2.02 (Cornuet & Luikart 1996), using the Wilcoxon test with 1000 iterations, and assuming both the infinite allele (IAM) and the stepwise mutation model (SMM). Tests for bottlenecks were conducted for the 1998 samples (in Ponds 1 and 3 with adults and larvae pooled) for loci in Hardy–Weinberg equilibrium.

On average 4% of adult newts from ponds where *T. cristatus* and *T. marmoratus* both occur are  $F_1$ -hybrids (Arntzen & Wallis 1991). In this study, adult hybrids were identified morphologically and larval hybrids were identified genetically with diagnostic microsatellites (Jehle *et al.* 2000), and excluded from the analysis. The frequency of introgressed protein loci is approximately 0.3% (Arntzen & Wallis 1991). For microsatellites, 0.7% of all individuals morphologically designated as either *cristatus* or *marmoratus* had an 'alien' allele in the species-diagnostic locus (see below). Such alleles were excluded from  $N_b$  and  $N_e$  calculations.

#### Calculations of $N_b$ and $N_e$

Variation in allele frequencies between adults and their larval progeny was used for calculating  $N_b$  (Ponds 1 and 3). On the assumption that our sampling was nondestructive,

we followed Plan I from Waples (1989a). Larval allele frequencies from 1986 and 1989 were compared with adult allele frequencies from 1998 for calculating  $N_e$  (Ponds 1 and 2), in this case following Plan II as larvae were removed (Waples 1989a). The standard variance of allele frequency change ( $F_c$ ) was calculated after Nei & Tajima (1981):

$$F_c = 1/K \sum (x_i - y_i) / [(x_i + y_i)^2 / (2 - x_i y_i)] \quad (1)$$

where  $K$  is the number of alleles, and  $i$  is the frequency of the respective allele at times  $x$  and  $y$ , respectively.  $F_c$  was then corrected for sample size and actual population size with Equations 12 ( $N_b$ ) and 11 ( $N_e$ ) from Waples (1989a):

$$N_b = t/2[F_c - 1/(2S_0) - 1/(2S_t) + 1/N] \quad (2)$$

$$N_e = t/2[F_c - 1/(2S_0) - 1/(2S_t)] \quad (3)$$

where  $t$  is the generation time,  $S_0$  and  $S_t$  represent sample sizes at times zero and time  $t$ , and  $N$  represents the adult population census size at time zero. In the study area, the mean number of reproductive years is 2.5 for *T. cristatus* and 4.0 for *T. marmoratus*, following the onset of sexual maturity at ages of three and four years, respectively (Francillon-Vieillot *et al.* 1989; Arntzen & Hedlund 1990). In the absence of data on age-dependent reproductive success, the generation time  $t$  was estimated as 4.3 years ( $= 3 + 2.5/2$ ) for *T. cristatus* and as 6.0 years ( $= 4 + 4/2$ ) for *T. marmoratus*. For  $N_b$  estimates,  $t$  was set at one (cf. Scribner *et al.* 1997). Due to the presence of null alleles (see below), and further missing data for example due to the omission of alien alleles, sample size varied among loci, and we used the harmonic mean of  $S_0$  and  $S_t$ . Under the assumption that  $N_e$  follows a  $\chi^2$ -distribution, we used equation 16 from Waples (1989a) in order to calculate 95% confidence intervals:

$$(1 - \alpha) \text{ confidence interval for } F_c = nF_c / (\chi^2_{\alpha/2}[n]), \\ nF_c / (\chi^2_{1 - \alpha/2}[n]) \quad (4)$$

where  $n$  is the number of degrees of freedom associated with  $F_c$  [ $n = \Sigma(\text{number of alleles} - 1)$ ] and  $\chi^2_{\alpha/2}[n]$  is the critical  $\alpha/2$  chi-square value for  $n$  degrees of freedom. The confidence limits obtained were used in place of  $F_c$  to determine the confidence interval of  $N_b$  and  $N_e$ .

The assumptions underlying the calculation are that sampling is at random, selection and mutation are negligible, and that there is no migration between populations and no population substructure. *Tcri36* failed to reliably amplify the small 1989 sample for *T. cristatus* in Pond 1 and was excluded from the analysis.  $N_b$  and  $N_e$  calculations were performed with all loci together, and separately only for loci in Hardy–Weinberg equilibrium. Pond 2 contained only few *T. cristatus* and this species was not analysed.

**Table 1** Characterization of the polymorphic microsatellite loci observed in the newts *Triturus cristatus* and *T. marmoratus*. *n*, number of alleles;  $H_E$ , expected heterozygosity;  $H_O$ , observed heterozygosity; *P*, probability level of test for Hardy–Weinberg equilibrium; n.s., non significant

Locus	Allele	Pond 1			Pond 2		Pond 3	
		1989 larvae	1998 adult	1998 larvae	1986 larvae	1998 adult	1998 adult	1998 larvae
<i>T. cristatus</i>		8	35	40	—	—	168	87
<i>Tcri13</i>	<i>n</i>	3	3	3			4	2
	$H_O$	0.500	0.429	0.550	—	—	0.196	0.184
	$H_E$	0.577	0.455	0.486	—	—	0.223	0.169
	<i>P</i>	n.s.	n.s.	n.s.	—	—	n.s.	n.s.
<i>Tcri27</i>	<i>n</i>	4	8	6			6	6
	$H_O$	0.375	0.571	0.600	—	—	0.613	0.448
	$H_E$	0.325	0.619	0.537	—	—	0.553	0.531
	<i>P</i>	n.s.	n.s.	n.s.	—	—	n.s.	n.s.
<i>Tcri29</i>	<i>n</i>	3	3	3			5	4
	$H_O$	0.375	0.629	0.450	—	—	0.536	0.460
	$H_E$	0.667	0.567	0.537	—	—	0.588	0.547
	<i>P</i>	n.s.	n.s.	n.s.	—	—	n.s.	n.s.
<i>Tcri32</i>	<i>n</i>	2	4	5			7	9
	$H_O$	0.125	0.400	0.600	—	—	0.476	0.701
	$H_E$	0.125	0.410	0.628	—	—	0.513	0.713
	<i>P</i>	—	n.s.	n.s.	—	—	n.s.	n.s.
<i>Tcri35</i>	<i>n</i>	3	4	3			5	5
	$H_O$	0.500	0.571	0.500	—	—	0.571	0.540
	$H_E$	0.548	0.635	0.467	—	—	0.552	0.551
	<i>P</i>	n.s.	n.s.	n.s.	—	—	n.s.	n.s.
<i>Tcri36</i>	<i>n</i>	—	2	3			3	4
	$H_O$	—	0.457	0.600	—	—	0.470	0.563
	$H_E$	—	0.464	0.506	—	—	0.463	0.502
	<i>P</i>	—	n.s.	n.s.	—	—	n.s.	n.s.
<i>Tcri46b</i>	<i>n</i>	5	5	7			6	5
	$H_O$	0.375	0.314	0.500	—	—	0.238	0.230
	$H_E$	0.725	0.612	0.689	—	—	0.518	0.495
	<i>P</i>	*	**	**	—	—	**	**
<i>Tcri43</i>	<i>n</i>	3	6	6			9	8
	$H_O$	0.250	0.486	0.775	—	—	0.655	0.724
	$H_E$	0.414	0.570	0.646	—	—	0.684	0.762
	<i>P</i>	n.s.	n.s.	n.s.	—	—	*	n.s.
<i>T. marmoratus</i>		23	66	79	90	43	27	7
<i>Tcri27</i>	<i>n</i>	2	5	5	6	4	3	3
	$H_O$	0.304	0.515	0.430	0.222	0.209	0.259	0.286
	$H_E$	0.324	0.476	0.401	0.211	0.229	0.259	0.272
	<i>P</i>	n.s.	*	n.s.	n.s.	n.s.	n.s.	n.s.
<i>Tcri32</i>	<i>n</i>	4	4	4	6	6	4	2
	$H_O$	0.130	0.379	0.367	0.500	0.372	0.407	0.439
	$H_E$	0.126	0.326	0.375	0.470	0.386	0.514	0.363
	<i>P</i>	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.	n.s.
<i>Tcri35</i>	<i>n</i>	4	5	5	6	5	4	4
	$H_O$	0.522	0.530	0.430	0.467	0.349	0.407	0.429
	$H_E$	0.436	0.561	0.427	0.688	0.437	0.507	0.571
	<i>P</i>	n.s.	n.s.	n.s.	**	n.s.	n.s.	n.s.
<i>Tcri46b</i>	<i>n</i>	3	6	6	5	6	5	2
	$H_O$	0.043	0.318	0.076	0.256	0.326	0.148	0.000
	$H_E$	0.211	0.427	0.241	0.288	0.348	0.079	0.254
	<i>P</i>	**	**	**	n.s.	n.s.	**	n.s.
<i>Tcri43</i>	<i>n</i>	3	9	5	6	2	6	3
	$H_O$	0.000	0.364	0.291	0.133	0.023	0.296	0.000
	$H_E$	0.174	0.546	0.540	0.317	0.074	0.515	0.408
	<i>P</i>	**	**	**	**	n.s.	**	n.s.

\* $P < 0.05$ , \*\* $P < 0.01$ .

<i>T. cristatus</i>	1, 1998A	1, 1998L	3, 1998A	3, 1998L
1, 1998A	—	<b>0.78</b>	0.08	0.30
1, 1998L	<b>0.12</b>	—	0.07	0.23
3, 1998A	0.80	0.09	—	<b>0.59</b>
3, 1998L	0.89	0.21	<b>1.00</b>	—

  

<i>T. marmoratus</i>	1, 1989	1, 1998A	1, 1998L	2, 1986A	2, 1998A	3, 1998A
1, 1989A	—	<b>0.07</b>	<b>0.07</b>	0.04*	0.03*	0.11
1, 1998A	<b>0.04*</b>	—	<b>0.18</b>	0.85	0.91	0.07
1, 1998L	<b>0.14</b>	<b>0.50</b>	—	0.18	0.23	0.48
2, 1996A	0.23	0.46	0.92	—	<b>0.13</b>	0.07
2, 1998A	0.50	0.14	0.50	<b>0.08</b>	—	0.49
3, 1998A	0.35	0.12	0.60	0.35	0.25	—

\*significant at  $P < 0.05$ .

## Results

Estimated adult population census sizes ( $N$ ) were remarkably stable between samples. Pond 1 was inhabited by  $76.5 \pm 25.6$  (1989), or  $72.5 \pm 18.8$  (1998) *Triturus cristatus*, and  $146.0 \pm 55.4$  (1989), or  $147.4 \pm 28.4$  (1998) *T. marmoratus*. Pond 2 was inhabited by  $101 \pm 14.1$  (1986), or  $108.0 \pm 31.0$  (1998) *T. marmoratus*. Pond 3 in 1998 contained  $585.6 \pm 78.3$  *T. cristatus* and  $57.0 \pm 32.6$  *T. marmoratus*.

The analysed microsatellites yielded from two to nine alleles per locus and population (Table 1, see the Appendix I for the allelic data). Five of the eight loci were polymorphic in *T. marmoratus*, with one (*Tcri32*) having alleles of species-diagnostic length. In *T. cristatus*, *Tcri46b* showed significant heterozygote deficiencies (Table 1), with an estimated frequency of null alleles of 0.29. In *T. marmoratus*, heterozygote deficiencies were observed at two of the five loci (*Tcri46b* and *Tcri43*), with null allele frequencies of 0.21 and 0.23, respectively. No significant linkage disequilibrium between loci was observed (pairwise comparisons,  $P$ -values  $> 0.05$ ). Assuming that the mutation rate of microsatellites is  $10^{-5}$  (Ellegren 1995) and using the mean size of the two temporal samples, the probability of a single mutation was 0.0024 for *T. cristatus* and 0.0021 for *T. marmoratus* in Pond 1, and 0.0044 in Pond 2. Under the assumption of the IAM a bottleneck in *T. cristatus* was suggested in both ponds ( $P < 0.01$  in both cases), whereas under the SMM the deviation from mutation-drift equilibrium was not significant ( $P > 0.05$  in both cases). For *T. marmoratus*, the Wilcoxon test was not significant for Ponds 1 and 2 (IAM and SMM:  $P > 0.05$ ), but significant under the IAM for Pond 3 ( $P < 0.01$ , SMM:  $P > 0.05$ ).

Allele frequencies were subject to change between temporal samples. Common alleles maintained their frequency ranks, but some alleles that were rare in one sample were not observed in the other sample. *T. marmoratus* in 1989

**Table 2**  $P$ -values obtained from Wilcoxon tests for the significance of differences in number of alleles (above diagonal) and observed heterozygosity ( $H_O$ , below diagonal) between temporal samples within populations (in bold) and between populations. Pond number is given first, followed by study year; A, adults; L, larvae. *Triturus cristatus* samples from Pond 1 and *T. marmoratus* larvae from Pond 3 were omitted due to  $n < 10$  ( $n > 23$  for included samples)

**Table 3**  $F_{ST}$  values between temporal and spatial samples of *Triturus cristatus* and *T. marmoratus*. Due to the small sample size ( $n < 8$ ) *T. cristatus* adults collected in 1989 from Pond 1 and *T. marmoratus* larvae from Pond 3 are not considered

	<i>T. cristatus</i>	<i>T. marmoratus</i>
Temporal samples		
Pond 1, adult-larvae 1998	0.0006	0.0139
Pond 1, 1989–98	—	0.0063
Pond 2, 1986–98	—	0.0704
Pond 3, adult-larvae 1998	0.0005	—
Spatial samples (adults only)		
Pond 1–Pond 2	—	0.1758
Pond 1–Pond 3	0.0455	0.0816
Pond 2–Pond 3	—	0.1313

(Pond 1) had a lower genetic diversity than the other samples, which is associated with the significant differences in allele numbers or  $H_O$  (Table 2). No significant differences were observed for other pairwise temporal or spatial comparisons.  $F_{ST}$  for parent–offspring cohort comparisons was at least one order of magnitude smaller than between samples from the same populations involving longer time intervals (Table 3). With the exception of *T. marmoratus* in Pond 2, spatial  $F_{ST}$  was higher than temporal  $F_{ST}$ .

$F_C$ , reflecting the variation in allele frequency changes, ranged between 0.04 and 0.18 (Table 4), and was larger when measured over longer time intervals (data from Pond 1).  $N_b$  and  $N_e$  represented a fraction of  $N$ , and the upper confidence limits reached  $N$  only for *T. marmoratus* in Pond 3 (estimated as infinity).  $N_b$  ranged between 9.2 and 101.8 newts,  $N_e$  ranged between 9.6 and 13.4 newts.  $N_b/N$  ranged between 0.10 and 0.19, and  $N_e/N$  ranged between 0.09 and 0.16.  $N_b/N$  and  $N_e/N$  in *T. cristatus* were always equal to or higher than in *T. marmoratus*. For

**Table 4** Variables used for the calculations of effective population size [ $F_c$ , standard variance in allele frequency change according to Nei & Tajima (1981);  $t$ , number of generations between samples for  $N_b$ ;  $S_0$  and  $S_1$ , sample size at time zero and 1, respectively]; number of breeding adults ( $N_b$ ) and effective population size ( $N_e$ ) with associated 95% confidence interval;  $N_b/N$ , respectively  $N_e/N$  values of syntopic *Triturus cristatus* and *T. marmoratus* populations. Data from Pond 2 refer to *T. marmoratus*. \*alternative measures of  $N_e$  and  $N_b$  calculated only from loci which are in Hardy–Weinberg equilibrium

	$F_c$	$t$	$S_0$	$S_1$	$N_b, N_e$	$N_b/N, N_e/N$	$N_b, N_e^*$
Pond 1							
<i>T. cristatus</i>							
$N_b$	0.050	1	162.2	72.4	14.1 (8.7–27.7)	0.19	9.3 (4.5–17.7)
$N_e$	0.189	2.18	61.4	73	12.2 (4.7–47.9)	0.16	11.1 (4.0–49.3)
<i>T. marmoratus</i>							
$N_b$	0.043	1	106	128	15.2 (7.8–27.9)	0.10	62.8 (15.9–1737.1)
$N_e$	0.104	1.50	26	110.8	13.4 (5.3–43.0)	0.09	50.6 (7.7–∞)
Pond 2							
$N_e$	0.107	2.00	148.3	65	9.6 (6.0–19.5)	0.09	10.5 (4.5–21.4)
Pond 3							
<i>T. cristatus</i>							
$N_b$	0.0133	1	294	161.5	101.8 (41.0–506.4)	0.17	89.3 (35.5–408.9)
<i>T. marmoratus</i>							
$N_b$	0.144	1	44	11.6	10.6 (3.6–∞)	0.17	∞ (3.5–∞)

*T. cristatus*, removing *Tcri46b* had little effect on  $N_b$  or  $N_e$ . When removing such loci for *T. marmoratus*, however, in Pond 1  $N_b$  and  $N_e$  increased about fourfold, and in Pond 3  $N_e$  could not be defined.

## Discussion

Temporal population genetic studies are scarce compared to studies on spatial population structure. Microsatellites have the technical advantage that PCR amplifications are possible from samples taken decades ago for purposes other than DNA analysis (for example from fish scales, Miller & Kapuscinski 1997; Nielsen *et al.* 1999), and have proven to be useful for assessing the temporal stability of populations (e.g. Taylor *et al.* 1994; Tessier & Bernatchez 1999). For the temporal method of calculating  $N_e$ , the mutation rates of microsatellites are such that they provide a high level of within-population polymorphism, while keeping the probability of mutations at an insignificant level. The codominant nature of microsatellite markers provides approximately twice the precision compared to markers displaying dominant gene expression (Jorde *et al.* 1999). However, multiallelic microsatellite data sets, such as ours, at present render a calculation of  $N_b$  or  $N_e$  using likelihood approaches computationally prohibitive (Williamson & Slatkin 1999), and do not increase the power of  $N_e$  calculations based on linkage disequilibrium (Hill 1981; Schwartz *et al.* 1998).

Microsatellites have to be treated with some caution, as the assumption of Mendelian inheritance may not always be met (Ardren *et al.* 1999). Deviations from Hardy–Weinberg

equilibrium were relatively frequent, especially in *Triturus marmoratus*. As observed for some of the loci, it is unlikely that deviations were caused by population substructuring. Moreover, heterozygotes were typed in both sexes, indicating that none of the loci are sex-linked. Deviations from Hardy–Weinberg equilibrium were associated with missing PCR products, and we concluded that null alleles were the cause. The frequent occurrence of null alleles in species for which the PCR primers were not originally designed has been reported previously (Paetkau & Strobeck 1995). Allele frequencies, and therefore the temporal method, should not be seriously affected when the relative number of amplification failures is similar across samples. The estimates of  $N_e$  for *T. marmoratus* with the loci that were out of Hardy–Weinberg proportions excluded, inevitably yielded a larger confidence interval.

### Spatial vs. temporal genetic differentiation

As recommended for microsatellite data and moderate sample sizes (Gaggiotti *et al.* 1999), we used  $F_{ST}$  based on Weir & Cockerham (1984) to measure genetic differentiation. Between Ponds 1 and 3  $F_{ST}$  was twice as large for *T. marmoratus* as for *T. cristatus*, but data from additional populations would be required to test whether this is a general pattern, and the possibility has to be considered that  $F_{ST}$  might be affected by the different number and performance of the used loci. The high temporal  $F_{ST}$  of *T. marmoratus* in Pond 1 cannot be explained by genetic drift, as Pond 2 had a similar  $N_e$  but a much lower  $F_{ST}$ , and was associated with a low degree of genetic diversity in 1989.

That spatial  $F_{ST}$  exceeds temporal  $F_{ST}$  in a microsatellite data set was also observed by Nielsen *et al.* (1999) for salmonid fish, with, however, different sampling regimes and study objectives.

#### $N_b$ and $N_e$ of *T. cristatus* and *T. marmoratus*

The habit of breeding in small ponds makes large-bodied newts amenable to measure  $N_b$  and  $N_e$ . Although part of our study is based on whole larvae collected prior to the routine use of PCR, tissue can be sampled nondestructively and in a straightforward way. To fit the assumption that sampling is at random with respect to family structure, adults and larvae were captured over the whole pond area. In another amphibian, larval growth but not survival was significantly correlated with microsatellite heterozygosity (Rowe *et al.* 1999). As larval sampling was not size-selective, there was no evidence for biased  $N_b$  and  $N_e$  due to selection at early life stages. At present there is no method available that calculates  $N_e$  taking migration into account. Ignoring constant migration leads to  $N_e$  estimates that are too high (Lehmann *et al.* 1998), whereas an episodic migration from sources with different allele frequencies would probably bias the estimate downward. Radio-tracking of 44 adult *T. cristatus* and *T. marmoratus* at Ponds 1 and 3 did not result in any indications for newts to disperse (Jehle & Arntzen 2000; Jehle 2000). Juveniles are frequently speculated to be the most important life stage for amphibian dispersal, but recent studies on *T. cristatus* suggest that their migration behaviour is similar to that of adults (by following their directional cues, Hayward *et al.* 2000), and indicate a high degree of pond fidelity (Cummins & Swan 2000). Nevertheless, over a sampling period of more than one decade, some bias of  $N_e$  due to migration cannot be completely ruled out.

The bias in  $N_e$  estimates caused by overlapping generations can be regarded as small when a time period covering more than one generation is sampled (Jorde & Ryman 1995). This is the case in our study, but the lack of age-specific reproductive rates led to an only rough estimate of generation time. However, body size and fecundity, and not body size and age, are usually correlated in amphibians (Duellman & Trueb 1986), suggesting that reproductive rates do not markedly differ between age classes. Hybridization is another potential source of bias. In the analysis of a deer hybrid zone with 11 diagnostic microsatellites, without any  $F_1$ -hybrids in the data set, up to 40% of all individuals carried introgressed alleles (Goodman *et al.* 1999). We excluded alien alleles for  $N_b$  and  $N_e$  calculations, but introgressed alleles from nondiagnostic loci with species-specific frequencies might have biased the  $N_b$  and  $N_e$  estimates slightly upwards, similar to the effect of constant migration. For discussions of further possible violations of assumptions of the temporal method see: Waples (1989a);

Husband & Barrett (1992); Scribner *et al.* (1997); Ingvarsson & Olsson (1997); Miller & Kapuscinski (1997); and Lehmann *et al.* (1998).

As fewer loci were used in *T. marmoratus*, the confidence limits for  $N_e$  estimates are mostly wider than for *T. cristatus*. Caution should be taken for  $N_e$  estimates based on a small number of samples (Waples 1989a), and the indeterminate  $N_b$  for *T. marmoratus* from Pond 3 (loci out of Hardy-Weinberg equilibrium excluded) was indeed associated with a low sample size and small number of loci. Pooling adults and larvae from 1998 would have increased  $S_t$  in Pond 1, but proved unnecessary and would have introduced a bias due to the merging of generational cohorts. It has to be noted, however, that the substantial confidence intervals of  $N$ ,  $N_b$  and  $N_e$  become multiplied in the  $N_b/N$  and  $N_e/N$  ratios, and lower their accuracy. Pooling rare alleles (frequencies < 0.05) only marginally changed the estimated  $N_e$  and  $N_b$  values (results not shown).

$N_b/N$  and  $N_e/N$  were relatively stable across populations, and in the same range as the mean  $N_e/N$  of 0.10 measured across 102 taxa (ranging from plants to humans, Frankham 1995b). That behavioural mechanisms and/or sperm competition cause only few males to reproduce successfully is probably more likely than frequent reproductive failures of females. In Pond 1, but not in Pond 3,  $N_b/N$  and  $N_e/N$  are twice as large for *T. cristatus* than for *T. marmoratus*. *T. marmoratus* has a higher fecundity and longevity than *T. cristatus* (Arntzen & Hedlund 1990), but the promiscuous mating system of newts in combination with internal fertilizations and the lack of parental care render it impossible to obtain precise measures of variance in reproductive success from life-history data. It has to be also noted that species-specific differences might be caused by the different performance and number of loci used. Vucetich *et al.* (1997) showed that a fluctuation in population size can lead to the discrepancy of  $N_e/N$  being higher in theoretical than in empirical estimates. The population sizes in Ponds 1 and 2 were stable between the sampling dates, and also within the sampling period there were no excessive fluctuations (J. W. Arntzen, unpublished data).  $N_b$  as measured from one reproductive period is identical to  $N_e$  only when the study organism has a single breeding cycle. However, large-bodied newts can visit their breeding ponds for up to at least five seasons (Ellinger & Jehle 1997), implying that the true  $N_e$  is probably larger than the measured  $N_b$ . From this view, our data confirm that  $N_e$  decreases with increasing sampling interval (Vucetich *et al.* 1997), despite a stable census size.

#### *A comparison between newts (this study) and toads (Scribner et al. 1997)*

That life-history parameters influence  $N_e/N$  is well documented (e.g. Nunney 1993, 1996), but empirical comparisons between related taxa with different reproductive

strategies have so far been lacking. Scribner *et al.* (1997) used other marker loci but the identical calculation method for calculating  $N_b/N$  in *Bufo bufo*, and revealed values in the order of 20 times lower (0.005–0.012) than this study on newts. It is unknown how multiple matings, which are common in newts (Halliday 1998) but rare or absent in toads, influence  $N_b$  and  $N_e$ . Two factors, however, probably account for a lower variance in reproductive success in newts. First, female toads are outnumbered by males (Arntzen 1999), whereas newts reproduce at an even sex ratio (Faber 1995). Second, female toads deposit whole egg strings in a very restricted pond area, exposing them to catastrophic events such as predation or desiccation, whereas female newts singly distribute their eggs over vast ponds areas, reducing the risk of complete reproductive failures. Assuming that  $N$  in *T. cristatus* is typically one order of magnitude smaller than for *B. bufo* (Halley *et al.* 1996) leads to the conclusion that their effective population sizes are of the same order. *B. bufo* has the ability to migrate over larger distances than large-bodied newts, enabling the colonization of more distant breeding ponds (Halley *et al.* 1996). Human alterations to the landscape mostly result in pond loss and increased between-pond distances, and large-bodied newts seem to be more affected than *B. bufo*.

#### Tests for recent bottlenecks in *T. cristatus* populations

Due to the rapid reduction in the number of alleles, gene diversity in bottlenecked populations is higher than expected under the mutation-drift equilibrium of constant population size (Cornuet & Luikart 1996). Under the IAM, but not under the SMM, we found evidence that bottlenecks occurred in *T. cristatus* populations. However, microsatellites exhibit an intermediate mutation mechanism, and to be statistically conservative only results under the SMM should be considered (Luikart & Cornuet 1998). Perhaps the local colonization took place too many generations ago, or the bottleneck is overshadowed by the expansion of the local *T. cristatus* populations as a whole, or by migration between ponds. Still, across all studied ponds, in concordance with our hypothesis the statistical evidence for bottlenecks in *T. cristatus* is slightly stronger than for *T. marmoratus*, although the species-specific difference is somewhat difficult to interpret as the performance and number of loci was not identical.

#### Implications for conservation

The relative stability of  $N_b$  and  $N_e$  in the present study should not be regarded as typical, as the populations were chosen due to their large size, and, a posteriori, due to their long-term persistence. Low  $N_e$  leads to a rapid loss of allelic diversity, but the small number of studied populations precludes a correlation of  $N_e$  and heterozygosity

(see also Funk *et al.* 1999). The effective population sizes of newts might be especially sensitive to suboptimal habitat quality, for example through the disruption of spermatophore transfer at turbid ponds with muddy substrates. Given that microsatellite heterozygosity may be correlated with fitness parameters (Coltman *et al.* 1998; Rowe *et al.* 1999), experimental studies to compare  $N_b$  or  $N_e$  under regimes of varying pond quality might provide important insights into whether there is greater genetic threat to amphibian populations under suboptimal conditions than in good-quality habitats, and whether such mechanisms could have contributed to the observed global amphibian population decline.

The typical amphibian population structure is conducive to high levels of inbreeding. Our study reveals  $N_e$  values for large-bodied newts that are larger than for other amphibians, but still one order of magnitude smaller than the actual census size. In metapopulations with relatively constant gene flow between subpopulations, the overall  $N_e$  is higher relative to a panmictic population living in the same resource, but falls below this level when demic extinction and recolonization rates are high (Whitlock & Barton 1997). European pond breeding amphibians often consist of populations with a census size of < 100 individuals. Stable metapopulations have been described for situations where ponds are to some degree interconnected (Rowe *et al.* 2000), but due to landscape fragmentation many populations are now restricted to completely isolated habitat remnants. The effective size of amphibian populations at the landscape scale could be alarmingly low, with loss of alleles and random genetic drift dominating the structural dynamics. Future studies could combine  $N_b$  or  $N_e$  measures with interdemic migration rates, and extrapolate such data sets over the landscape level in order to assess whether from a genetic point of view amphibian populations in fragmented habitats are viable in the long term, and to what extent they require active management strategies.

#### Acknowledgements

This study was supported by the Austrian Science Foundation (FWF grant P-11852 BIO) and the NERC (GR3/10333). Samples were collected under the permits 97/204 and 98/226/AUT of the Ministère de l'Environnement, Paris. We thank Dominique Bardou for hospitality. Doris Bachtrog, Gordon Luikart, Kim Scribner and two anonymous reviewers provided constructive comments on the manuscript.

#### References

- Ardren WR, Borer S, Thrower F, Joyce JE, Kapuscinski AR (1999) Inheritance of 12 microsatellite loci in *Oncorhynchus mykiss*. *Journal of Heredity*, **90**, 529–536.
- Arntzen JW (1999) Sexual selection and male mate choice in the common toad, *Bufo bufo*. *Ethology, Ecology and Evolution*, **11**, 407–414.

- Arntzen JW, Hedlund L (1990) Fecundity of the newts *Triturus cristatus*, *T. marmoratus* and their natural hybrids in relation to species coexistence. *Holarctic Ecology*, **13**, 325–332.
- Arntzen JW, Wallis G (1991) Restricted gene flow in a moving hybrid zone of newts (*Triturus cristatus* and *T. marmoratus*) in western France. *Evolution*, **45**, 805–826.
- Begon M (1979) *Investigating Animal Abundance: Capture-Recapture for Biologists*. Edward Arnold, London.
- Berven KA, Grudzien TA (1990) Dispersal in the wood frog (*Rana sylvatica*): implications for genetic population structure. *Evolution*, **44**, 2047–2056.
- Coltman DW, Bowen WD, Wright JM (1998) Birth weight and neonatal survival of harbour seal pups are positively correlated with genetic variation measured with microsatellites. *Proceedings of the Royal Society of London B*, **265**, 803–809.
- Cornuet JM, Luikart G (1996) Description and power analysis of two tests for detecting recent population bottlenecks from allele frequency data. *Genetics*, **144**, 2001–2014.
- Crandall KA, Posada D, Vasco D (1999) Effective population sizes: missing measures and missing concepts. *Animal Conservation*, **2**, 317–320.
- Cummins CA, Swan MJS (2000) Long-term survival and growth of free-living great crested newts (*Triturus cristatus*) PIT-tagged at metamorphosis. *Herpetological Journal*, **10**, 177–182.
- Driscoll DA (1999) Genetic neighbourhood and effective population size for two endangered frogs. *Biological Conservation*, **88**, 221–229.
- Duellman W, Trueb L (1986) *Biology of Amphibians*. McGraw-Hill, New York.
- Easteal S (1985) The ecological genetics of introduced populations of the giant toad *Bufo marinus* II. Effective Population size. *Genetics*, **110**, 107–122.
- Ellegren H (1995) Mutation rates at porcine microsatellite loci. *Mammalian Genome*, **6**, 376–377.
- Ellinger N, Jehle R (1997) Struktur und Dynamik einer Donaukammolch-Population (*Triturus dobrogicus* Kirtzescu 1903) am Endelteich bei Wien: Ein Überblick über neun Untersuchungsjahre. In: *Populationsbiologie von Amphibien: eine Langzeitstudie auf der Wiener Donauinsel* (eds Hödl W, Jehle R, Gollmann G), pp. 133–150. Stapfia 51, Linz.
- Faber H (1995) Saisonale Dynamik der Geschlechterrelation beim Bergmolch, *Triturus alpestris alpestris* (LAURENTI, 1768), im aquatischen Lebensraum. *Herpetozoa*, **8**, 117–124.
- Fiumera AC, Parker PG, Fuerst PA (2000) Effective population size and maintenance of genetic diversity in captive-bred populations of a lake Victoria cichlid. *Conservation Biology*, **14**, 886–892.
- Francillon-Vieillot H, Arntzen JW, Géraudie J (1989) Age, growth and longevity of sympatric *Triturus cristatus*, *T. marmoratus* and their hybrids (Amphibia, Urodela): a skeletochronological comparison. *Journal of Herpetology*, **24**, 13–22.
- Frankham R (1995a) Conservation genetics. *Annual Reviews in Genetics*, **29**, 305–327.
- Frankham R (1995b) Effective population size/adult population size ratios in wildlife: a review. *Genetical Research, Cambridge*, **66**, 95–107.
- Funk WC, Tallmon DA, Allendorf FW (1999) Small effective population size in the long-toed salamander. *Molecular Ecology*, **8**, 1633–1640.
- Gaggiotti OE, Lange O, Rassmann K, Gliddon C (1999) A comparison of two indirect methods for estimating average levels of gene flow using microsatellite data. *Molecular Ecology*, **8**, 1513–1520.
- Gill DE (1978) Effective population size and interdemographic migration rates in a metapopulation of the red-spotted newt, *Notophthalmus viridescens* (Rafinesque). *Evolution*, **32**, 839–849.
- Goodman SJ, Barton NB, Swanson G, Abernethy K, Pemberton JM (1999) Introgression through rare hybridisation: a genetic study of a hybrid zone between red and sika deer (genus *Cervus*) in Argyll, Scotland. *Genetics*, **152**, 355–371.
- Guo SW, Thompson EA (1992) Performing the exact test of Hardy-Weinberg proportions for multiple alleles. *Biometrics*, **48**, 361–362.
- Haberl M, Tautz D (1999) Comparative allele sizing can produce inaccurate allele size differences for microsatellites. *Molecular Ecology*, **8**, 1347–1350.
- Halley J, Oldham RS, Arntzen JW (1996) Predicting the persistence of amphibian populations with the help of a spatial model. *Journal of Applied Ecology*, **33**, 455–470.
- Halliday T (1998) Sperm competition in amphibians. In: *Sperm Competition and Sexual Selection* (eds Birkhead TR, Møller AP), pp. 465–502. Academic Press, London.
- Hayward R, Oldham RS, Watt PJ, Head SM (2000) Dispersion patterns of young great crested newts (*Triturus cristatus*). *Herpetological Journal*, **10**, 129–136.
- Hedrick PW (1999) Perspective: Highly variable loci and their interpretation in evolution and conservation. *Evolution*, **53**, 313–318.
- Hill W (1981) Estimation of effective population size from data on linkage disequilibrium. *Genetical Research, Cambridge*, **38**, 209–261.
- Husband BC, Barrett SCH (1992) Effective population size and genetic drift in tristylous *Eichhornia paniculata* (Pontederiaceae). *Evolution*, **46**, 1875–1890.
- Ingvarsson PK, Olsson K (1997) Hierarchical genetic structure and effective population size in *Phalacrocorax substriatus*. *Heredity*, **79**, 153–161.
- Jehle R (2000) The terrestrial summer habitat of radio-tracked crested and marbled newts (*Triturus cristatus* and *T. marmoratus*). *Herpetological Journal*, **10**, 137–142.
- Jehle R, Arntzen JW (2000) Post-breeding migrations of newts (*Triturus cristatus* and *T. marmoratus*) with contrasting ecological requirements. *Journal of Zoology, London*, **251**, 297–306.
- Jehle R, Bouma P, Sztatecsny M, Arntzen JW (2000) High aquatic niche overlap in the newts *Triturus cristatus* and *T. marmoratus* (Amphibia, Urodela). *Hydrobiologia*, **437**, 149–155.
- Jorde PE, Palm S, Ryman N (1999) Estimating genetic drift and effective population size from temporal shifts in dominant gene marker frequencies. *Molecular Ecology*, **8**, 1171–1178.
- Jorde PE, Ryman N (1995) Temporal allele frequency change and estimation of effective size in populations with overlapping generations. *Genetics*, **139**, 1077–1090.
- Lehmann T, Hawley WA, Grebert H, Collins FH (1998) The effective population size of *Anopheles gambiae* in Kenya: Implications for population structure. *Molecular Biology and Evolution*, **15**, 264–276.
- Luikart G, Cornuet JM (1998) Empirical evaluation of a test for identifying recently bottlenecked populations from allele frequency data. *Conservation Biology*, **12**, 228–237.
- Luikart G, England PR (1999) Statistical analysis of microsatellite DNA data. *Trends in Ecology and Evolution*, **14**, 253–256.
- Marshall TC, Slate J, Kruuk LEB, Pemberton JM (1998) Statistical confidence for likelihood-based paternity inference in natural populations. *Molecular Ecology*, **7**, 639–655.
- Merrell DJ (1968) A comparison of the 'effective size' of breeding populations of the leopard frog, *Rana pipiens*. *Evolution*, **22**, 274–283.

- Miller LM, Kapuscinski AR (1997) Historical analysis of genetic variation reveals low effective population size in a northern pike (*Esox lucius*) population. *Genetics*, **147**, 1249–1258.
- Nei M, Tajima F (1981) Genetic drift and estimation of effective population size. *Genetics*, **98**, 625–640.
- Newman D, Pilson D (1997) Increased probability of extinction due to decreased effective population size: experimental populations of *Clarkia pulchella*. *Evolution*, **51**, 354–362.
- Nielsen EE, Hansen MM, Loeschke V (1999) Genetic variation in time and space: Microsatellite analysis of extinct and extant populations of Atlantic salmon. *Evolution*, **53**, 261–268.
- Nunney L (1993) The influence of mating system and overlapping generations on effective population size. *Evolution*, **47**, 1329–1341.
- Nunney L (1996) The influence of variation in female fecundity on effective population size. *Biological Journal of the Linnean Society*, **59**, 411–425.
- Paetkau D, Strobeck C (1995) The molecular basis and evolutionary history of a microsatellite null allele in bears. *Molecular Ecology*, **4**, 519–520.
- Raymond M, Rousset F (1995) GENEPOP (Version 2.1): population genetic software for exact test and ecumenism. *Journal of Heredity*, **86**, 248–249.
- Rowe G, Beebee TJC, Burke T (1999) Microsatellite heterozygosity, fitness and demography in natterjack toads *Bufo calamita*. *Animal Conservation*, **2**, 85–92.
- Rowe G, Beebee TJC, Burke T (2000) A microsatellite analysis of natterjack toad, *Bufo calamita*, metapopulations. *Oikos*, **88**, 641–651.
- Sambrook J, Fritsch EF, Maniatis T (1989) *Molecular Cloning: a Laboratory Manual*. 2nd edn. Cold Spring Harbour Laboratory Press, New York.
- Schwartz MK, Tallmon DA, Luikart G (1998) Review of DNA-based census and effective population size estimators. *Animal Conservation*, **1**, 293–299.
- Scribner KT, Arntzen JW, Burke T (1997) Effective number of breeding adults in *Bufo bufo* estimated from age-specific variation at minisatellite loci. *Molecular Ecology*, **6**, 701–712.
- Seppä P, Laurila A (1999) Genetic structure of island populations of the anurans *Rana temporaria* and *Bufo bufo*. *Heredity*, **82**, 309–317.
- Taylor AC, Sherwin WB, Wayne RK (1994) Genetic variation in a bottlenecked species: the northern hairy-nosed wombat *Larsiohinus kreftii*. *Molecular Ecology*, **3**, 277–290.
- Tessier N, Bernatchez L (1999) Stability of population structure and genetic diversity across generations assessed by microsatellites among sympatric populations of landlocked Atlantic salmon (*Salmo salar* L.). *Molecular Ecology*, **8**, 169–179.
- Vucetich JA, Waite TA, Nunney L (1997) Fluctuating population size and the ratio of effective to census population size. *Evolution*, **51**, 2017–2021.
- Waples RS (1989a) A generalised approach for estimating effective population size from temporal change in gene frequency. *Genetics*, **121**, 379–391.
- Waples RS (1989b) Temporal variation in allele frequencies: testing the right hypothesis. *Evolution*, **43**, 1236–1251.
- Weir BS, Cockerham CC (1984) Estimating *F*-statistics for the analysis of population structure. *Evolution*, **38**, 1358–1370.
- Whitlock MC, Barton NH (1997) The effective size of a subdivided population. *Genetics*, **146**, 427–441.
- Williamson EG, Slatkin M (1999) Using maximum likelihood to estimate population size from temporal changes in allele frequencies. *Genetics*, **152**, 755–761.

---

This study stems from the collaboration of several Institutions and persons interested in a wide range of ecological and evolutionary questions in amphibians. It is part of Robert Jehle's completed PhD research on the population ecology of European large-bodied newts, supervised by Walter Hödland Pim Arntzen. The laboratory work was conducted in the molecular ecology laboratory of Terry Burke, where Andy Krupa is a research technician.

---

## Appendix I

Allele frequency data for eight polymorphic microsatellite data observed in the newts *Triturus cristatus* and *T. marmoratus*. *Tcri32* is diagnostic between the species. For *Tcri27*, *Tcri35*, and *Tcri46b*, one or more alleles occur in one species but not in the other. *n* = per-locus sample size

Locus	Allele	Pond 1			Pond 2		Pond 3	
		1989 larvae	1998 adult	1998 larvae	1986 larvae	1998 adult	1998 adult	1998 larvae
<i>T. cristatus</i>								
<i>Tcri13</i>	A	0.429	0.227	0.308	—	—	0.129	0.093
	B	0.143	0.091	0.051	—	—	0.003	0.000
	C	0.429	0.682	0.641	—	—	0.858	0.907
	D	0.000	0.000	0.000	—	—	0.010	0.000
	<i>n</i>	14	66	78	—	—	302	172
<i>Tcri27</i>	A	0.333	0.015	0.194	—	—	0.000	0.000
	B	0.000	0.030	0.000	—	—	0.034	0.021
	C	0.333	0.545	0.597	—	—	0.541	0.486
	D	0.000	0.015	0.000	—	—	0.021	0.039
	E	0.000	0.152	0.097	—	—	0.219	0.277
	F	0.167	0.015	0.014	—	—	0.014	0.007
	G	0.000	0.167	0.083	—	—	0.000	0.000
	H	0.167	0.061	0.014	—	—	0.171	0.170
<i>n</i>	16	66	72	—	—	282	146	
<i>Tcri29</i>	A	0.250	0.132	0.243	—	—	0.169	0.205
	B	0.000	0.000	0.000	—	—	0.003	0.000
	C	0.250	0.309	0.176	—	—	0.325	0.295
	D	0.500	0.559	0.581	—	—	0.460	0.473
	E	0.000	0.000	0.000	—	—	0.043	0.027
	<i>n</i>	16	68	74	—	—	302	146
<i>Tcri32</i>	A	0.000	0.000	0.000	—	—	0.004	0.030
	B	0.000	0.222	0.229	—	—	0.453	0.375
	C	0.000	0.056	0.010	—	—	0.020	0.024
	D	0.000	0.000	0.019	—	—	0.059	0.083
	E	0.000	0.000	0.000	—	—	0.000	0.006
	F	0.000	0.000	0.000	—	—	0.000	0.006
	G	0.75	0.500	0.514	—	—	0.313	0.298
	H	0.25	0.222	0.229	—	—	0.148	0.167
	I	0.000	0.000	0.000	—	—	0.004	0.006
	<i>n</i>	14	44	80	—	—	256	168
<i>Tcri35</i>	A	0.000	0.000	0.000	—	—	0.007	0.026
	B	0.214	0.333	0.444	—	—	0.217	0.208
	C	0.000	0.030	0.074	—	—	0.086	0.110
	D	0.571	0.424	0.278	—	—	0.569	0.565
	E	0.214	0.212	0.000	—	—	0.121	0.101
	<i>n</i>	14	66	54	—	—	304	154
<i>Tcri36</i>	A	—	0.000	0.000	—	—	0.007	0.012
	B	—	0.500	0.513	—	—	0.493	0.476
	C	—	0.500	0.474	—	—	0.500	0.494
	D	—	0.000	0.013	—	—	0.000	0.018
	<i>n</i>	—	64	78	—	—	306	164
<i>Tcri46b</i>	A	0.000	0.000	0.014	—	—	0.021	0.012
	B	0.063	0.000	0.054	—	—	0.080	0.102
	C	0.483	0.065	0.054	—	—	0.000	0.000
	D	0.25	0.194	0.122	—	—	0.003	0.012
	E	0.25	0.468	0.378	—	—	0.626	0.651
	F	0.000	0.242	0.311	—	—	0.264	0.223
	G	0.063	0.032	0.068	—	—	0.006	0.000
	<i>n</i>	16	62	74	—	—	326	166

## Appendix I continued

Locus	Allele	Pond 1			Pond 2		Pond 3	
		1989 larvae	1998 adult	1998 larvae	1986 larvae	1998 adult	1998 adult	1998 larvae
<i>Tcri43</i>	A	0.000	0.000	0.000	—	—	0.019	0.065
	B	0.071	0.034	0.013	—	—	0.036	0.018
	C	0.000	0.121	0.118	—	—	0.062	0.101
	D	0.000	0.000	0.000	—	—	0.023	0.048
	E	0.714	0.431	0.382	—	—	0.179	0.185
	F	0.214	0.345	0.408	—	—	0.308	0.238
	G	0.000	0.017	0.039	—	—	0.351	0.327
	H	0.000	0.052	0.039	—	—	0.016	0.000
	I	0.000	0.000	0.000	—	—	0.006	0.018
<i>n</i>	14	58	76	—	—	308	168	
<i>T. marmoratus</i>								
<i>Tcri27</i>	B	0.000	0.009	0.000	0.000	0.000	0.000	0.000
	C	0.000	0.018	0.009	0.014	0.030	0.000	0.083
	D	0.000	0.000	0.000	0.007	0.000	0.000	0.000
	E	0.656	0.554	0.491	0.076	0.091	0.140	0.083
	F	0.344	0.366	0.466	0.854	0.833	0.840	0.833
	G	0.000	0.054	0.026	0.042	0.045	0.020	0.000
	H	0.000	0.000	0.009	0.007	0.000	0.000	0.000
	<i>n</i>	32	112	116	144	66	50	12
<i>Tcri32</i>	J	0.026	0.000	0.000	0.011	0.036	0.019	0.000
	K	0.053	0.032	0.026	0.052	0.036	0.000	0.000
	L	0.921	0.079	0.083	0.155	0.036	0.185	0.000
	M	0.000	0.802	0.776	0.695	0.774	0.667	0.786
	N	0.026	0.087	0.115	0.080	0.071	0.130	0.214
	O	0.000	0.000	0.000	0.006	0.000	0.000	0.000
	P	0.000	0.000	0.000	0.000	0.048	0.000	0.000
	<i>n</i>	38	126	156	174	84	54	14
<i>Tcri35</i>	A	0.000	0.000	0.000	0.185	0.000	0.000	0.000
	B	0.028	0.016	0.008	0.012	0.025	0.000	0.000
	C	0.000	0.000	0.008	0.238	0.013	0.019	0.071
	D	0.278	0.306	0.249	0.393	0.025	0.222	0.214
	E	0.611	0.564	0.660	0.155	0.688	0.667	0.643
	F	0.083	0.105	0.076	0.000	0.250	0.093	0.071
	G	0.000	0.008	0.000	0.018	0.000	0.000	0.000
	<i>n</i>	36	124	132	168	80	54	14
<i>Tcri46b</i>	B	0.000	0.012	0.021	0.007	0.027	0.206	0.200
	C	0.357	0.023	0.042	0.074	0.041	0.088	0.000
	D	0.214	0.233	0.042	0.797	0.770	0.059	0.000
	E	0.429	0.512	0.760	0.101	0.108	0.471	0.800
	F	0.000	0.186	0.115	0.020	0.027	0.176	0.000
	G	0.000	0.035	0.021	0.000	0.027	0.000	0.000
	<i>n</i>	14	86	96	148	74	34	10
<i>Tcri43</i>	A	0.000	0.028	0.029	0.000	0.000	0.000	0.000
	B	0.200	0.009	0.000	0.043	0.000	0.025	0.000
	C	0.200	0.019	0.065	0.054	0.000	0.100	0.000
	D	0.000	0.123	0.065	0.098	0.000	0.075	0.000
	E	0.400	0.500	0.529	0.565	0.417	0.500	0.500
	F	0.200	0.236	0.312	0.228	0.583	0.225	0.250
	G	0.000	0.009	0.000	0.011	0.000	0.075	0.250
	H	0.000	0.066	0.000	0.000	0.000	0.000	0.000
	I	0.000	0.009	0.000	0.000	0.000	0.000	0.000
	<i>n</i>	10	106	138	92	12	40	8