

RESEARCH ARTICLE

# Optional Endoreplication and Selective Elimination of Parental Genomes during Oogenesis in Diploid and Triploid Hybrid European Water Frogs

Dmitry Dedukh<sup>1</sup>, Spartak Litvinchuk<sup>2</sup>, Juriy Rosanov<sup>2</sup>, Glib Mazepa<sup>3,4</sup>, Alsu Saifitdinova<sup>1</sup>, Dmitry Shabanov<sup>4</sup>, Alla Krasikova<sup>1\*</sup>

**1** Saint-Petersburg State University, Saint-Petersburg, Russia, **2** Institute of Cytology, Russian Academy of Sciences, Saint-Petersburg, Russia, **3** Department of Ecology and Genetic, Evolutionary Biology Centre, Uppsala University, Uppsala, Sweden, **4** V.N. Karazin Kharkiv National University, Kharkiv, Ukraine

\* [alla.krasikova@gmail.com](mailto:alla.krasikova@gmail.com)



OPEN ACCESS

**Citation:** Dedukh D, Litvinchuk S, Rosanov J, Mazepa G, Saifitdinova A, Shabanov D, et al. (2015) Optional Endoreplication and Selective Elimination of Parental Genomes during Oogenesis in Diploid and Triploid Hybrid European Water Frogs. PLoS ONE 10(4): e0123304. doi:10.1371/journal.pone.0123304

**Academic Editor:** Pawel Michalak, Virginia Tech Virginia, UNITED STATES

**Received:** November 21, 2014

**Accepted:** March 3, 2015

**Published:** April 20, 2015

**Copyright:** © 2015 Dedukh et al. This is an open access article distributed under the terms of the [Creative Commons Attribution License](http://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

**Data Availability Statement:** All relevant data are within the paper and its Supporting Information files.

**Funding:** This work was funded in part by the Russian Foundation for Basic Research (<http://www.rfbr.ru/rffi/eng>) grant 12-04-01277 to SL and JR and by the grants of the President of Russia (<https://grants.extech.ru/>) NS-3553.2014.4 to AK, DD and AS and JC-3609.2014.4 to AK and DD. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

## Abstract

Incompatibilities between parental genomes decrease viability of interspecific hybrids; however, deviations from canonical gametogenesis such as genome endoreplication and elimination can rescue hybrid organisms. To evaluate frequency and regularity of genome elimination and endoreplication during gametogenesis in hybrid animals with different ploidy, we examined genome composition in oocytes of di- and triploid hybrid frogs of the *Pelophylax esculentus* complex. Obtained results allowed us to suggest that during oogenesis the endoreplication involves all genomes occurring before the selective genome elimination. We accepted the hypothesis that only elimination of one copied genome occurs pre-meiotically in most of triploid hybrid females. At the same time, we rejected the hypothesis stating that the genome of parental species hybrid frogs co-exist with is always eliminated during oogenesis in diploid hybrids. Diploid hybrid frogs demonstrate an enlarged frequency of deviations in oogenesis comparatively to triploid hybrids. Typical for hybrid frogs deviations in gametogenesis increase variability of produced gametes and provide a mechanism for appearance of different forms of hybrids.

## Introduction

Interspecific hybridization usually leads to death or sterility of hybrid animals [1], [2]. However, in vertebrate hybrids, deviations from canonical gametogenesis and meiosis give rise to a variety of successful reproduction modes, such as parthenogenesis, gynogenesis, kleptogenesis and hybridogenesis [3–7]. Such deviations in gametogenesis include chromosomal endoreplication and elimination [8–9]. Endoreplication in germ cells leads to the formation of gametes with a multiple increase of chromosomal number [10–12]. Chromatin elimination occurring in germ cells leads to selective (in case of hybridogenesis and pre-equalizing hybrid meiosis) or nonselective (in case

**Competing Interests:** The authors have declared that no competing interests exist.

of kleptogenesis) deletion of genome part in gametes [5], [7], [13–15]. It remains unclear how these processes are realized during gametogenesis in hybrid animals. In particular, frequency and accuracy of genome elimination and endoreplication are poorly investigated.

Both chromosomal elimination and endoreplication take place during gametogenesis in hybrid European water frogs of the *Pelophylax esculentus* complex, which represents a widely used model for studying interspecies hybridization [14], [16–19]. This complex consists of two parental species, the lake frog (*P. ridibundus*, RR genotype,  $2n = 26$ ) and the pool frog (*P. lessonae*, LL genotype,  $2n = 26$ ), and their natural hybridogenetic form—the edible frog (*P. esculentus*,  $2n = 26$ ) with RL genotype [16], [17].

Not only diploid, but also two forms of triploid hybrids (with LLR and RRL genotypes,  $3n = 39$ ) exist in different *P. esculentus* population systems [6], [20]. *P. esculentus* triploids are especially abundant in the Seversky Donets river basin (Eastern Ukraine) [21–23]. Indeed, water frogs population systems found in Kharkiv region of Eastern Ukraine are represented by *P. ridibundus* species (R type), pure hybrid population systems (E type), population systems of R-E type where *P. ridibundus* co-exists with hybrids, and rare population systems inhabited by both parental and hybrid species (R-L-E type) [21].

In central European population systems, where *P. esculentus* usually co-exists with *P. lessonae* (L-E type), diploid hybrid frogs produce gametes with genome of *P. ridibundus* (R genome) [18], [19]. Previous studies of rare R-E population systems in Central Europe suggested that in diploid hybrid frogs, R genome is eliminated premeiotically while L genome is transmitted into gametes to produce hybrid frogs after crossing with parental species [6], [20]. It was thus proposed that type of transmitted genome is complementary to genome of parental species co-existing with hybrid frogs [6], [20]. We asked whether this regularity is true for *P. esculentus* population systems from Eastern Ukraine. We also hypothesized that in diploid hybrid frogs, endoreplication should occur during gametogenesis to produce diploid female gametes. Importantly, the diversity of chromosomal sets in gametes produced by triploid *P. esculentus* females has not been studied so far. Likewise, it is unknown whether chromosomes are eliminated and/or endoreplicated within the germ line in triploid frogs. Our additional aim was to check whether in triploid hybrid females single copied genome is eliminated premeiotically while double copied genome forms bivalents.

Cytogenetic analyses of germ cell karyotype in hybrid frogs reveals the changes in chromosomal number in gametogenesis, as well as the origin of diploid and triploid hybrids in different population systems. In amphibian females, parental chromosomes identification is possible by the analysis of giant lampbrush chromosomes (LBCs) obtained from growing oocytes [10], [19], [24], [25].

Lampbrush chromosomes from European water frogs were characterized in 1972 [26]. However reliable species identification was not performed and it was unclear whether analyzed frogs referred to *P. esculentus* complex [6], [27]. In 1979 Graf and Müller described lampbrush chromosomes from *P. esculentus* [28]. Precise identification of parental chromosomes in oocytes of hybrid animals was impossible until 1990 when Bucci and coauthors characterized LBCs of *P. ridibundus* and *P. lessonae* from Poland and pointed out dissimilarities between LBCs of parental species [19]. Analyzing morphological resemblance with LBCs of parental species the authors were able to identify chromosomes in oocytes of hybrid animals.

To test our hypotheses we examined lampbrush chromosome sets in oocytes from diploid and triploid hybrid *P. esculentus* frogs taken from the population systems of R-E type located in the East of the Ukraine. We found unusual chromosomal sets in growing oocytes connected with hybridogenetic way of reproduction. Mechanisms which lead to formation of oocytes with unusual chromosomal sets and contribution of female gametes to the maintenance of R-E hybrid *P. esculentus* population systems are discussed.

## Materials and Methods

### Samples studied

The European water frogs were sampled in the Kharkiv and Donetsk regions (Eastern Ukraine). *P. ridibundus* (N = 3) and *P. lessonae* (N = 2) individuals were collected from the Dnieper River basin in Krasnokutsk district proximate to hybrid formation centers. Hybrid females were taken from the Seversky Donets River basin. 13 triploid hybrid females with RRL genotype, 5 triploid hybrid females with LLR genotype and 9 diploid females with RL genotype were gathered from the population system of R-E type (S1 Table). All manipulations with animals were carried out in accordance with the national and international guidelines. The field studies did not involve endangered or protected species. Collected specimens are not listed in IUCN Redlist or by CITES. All specimens were collected in the regions of Ukraine, which are not considered as protected areas, thus no specific permissions were required for these locations. Techniques used to capture, tissue sampling and euthanasia sought to minimize animal suffering and were in accordance with recommendations of the Herpetological Animal Care and Use Committee (HACC) of the American Society of Ichthyologists and Herpetologists (available at: <http://www.asih.org/publications>). Each individual was anaesthetized by methoxyethane or submersion in a 1% solution of 3-aminobenzoic acid ethyl ester (MS 222). All procedures were approved by the Scientific Committee of the Biology Department of Saint-Petersburg State University.

### DNA flow cytometry

Genome composition of all frogs was established by measurement the DNA amount per nucleus using flow cytometer constructed at the Institute of Cytology, Russian Academy of Sciences, St. Petersburg. All animals were anesthetized MS222 1.5 g/l (Sigma) to take blood from the femoral vein. 0.1% Triton X100, 20 µg/ml ethidium bromide and 15 mM MgCl<sub>2</sub> were added to blood samples. Blood of grass frog (*Rana temporaria* Linnaeus, 1758) and male domestic mouse (*Mus musculus*; spleenocytes, C57B1 line) were used as reference standards as published previously [29], [30]. DNA histograms were created using the formula: DNA content = (samples mean peak)/(reference standard peak) × (reference standard genome size).

### Preparation of mitotic metaphase chromosomes

Mitotic metaphase chromosomes were obtained from intestine of parental species and hybrid frogs using standard manipulations. Intestinal tissue was dissected after injection of additional animals of both parental species and hybrids with 0.2–0.5 ml of a 0.3% solution of colchicine. Intestine was incubated in 0.05 M KCl for 20 minutes, then fixed in 3:1 ethanol-glacial acetic acid, where it was stored until slide preparation. Prior to metaphase plates preparation, intestine fragment was placed into a drop of 60% glacial acetic acid for 5 min and crushed. The cell suspension was dropped onto specimen slides previously heated to 60°C.

### Lampbrush chromosomes isolation

Lampbrush chromosomes were microsurgically isolated from *P. esculentus* oocytes according to standard procedure [31]. All females used in lampbrush chromosome analysis were not injected by colchicine or hormonal drugs. Prior to ovary isolation, frogs were anaesthetized with MS222 1.5 g/l (Sigma). Pieces of ovary were cut off from females and kept in the OR2 saline (82.5 mM NaCl, 2.5 mM KCl, 1 mM MgCl<sub>2</sub>, 1 mM CaCl<sub>2</sub>, 1 mM Na<sub>2</sub>HPO<sub>4</sub>, 5 mM HEPES (4-(2-hydroxyethyl)-1-piperazineethanesulfonic acid); pH 7.4). Nuclei were isolated from oocytes in the isolation medium “5:1” (83 mM KCl, 17 mM NaCl, 6.5 mM Na<sub>2</sub>HPO<sub>4</sub>, 3.5 mM

$\text{KH}_2\text{PO}_4$ , 1 mM  $\text{MgCl}_2$ , 1 mM DTT (dithiothreitol); pH 7.0–7.2) by jeweler forceps under the observation at Leica MZ16 stereomicroscope. Each nucleus was transferred into chamber attached to a specimen slide filled with one-fourth strength “5:1” medium with the addition of 0.1% paraformaldehyde and 0.01% 1 M  $\text{MgCl}_2$  where nuclear envelopes were removed. Then slide preparations were centrifuged for 30 min at +4°C, 4000 rpm, fixed in 2% paraformaldehyde in 1x phosphate buffered saline (PBS) for 30 min at RT, and post-fixed in 50% ethanol for 5 min and 70% ethanol overnight (at +4°C). Preparations were not dried before immunostaining but were dehydrated in 96% ethanol for 5 min and air dried before cytological observation or FISH procedures.

### Fluorescence *in situ* hybridization

FISH with telomeric probe was carried out on lampbrush and metaphase chromosomes as described previously [32]. Metaphase plates were pre-treated with RNase A (100–200  $\mu\text{g}/\text{ml}$ ) for 1 h, pepsin (0.01% in 0.01 N HCl) for 10 min and then post-fixed in formaldehyde (1% in PBS, 50 mM  $\text{MgCl}_2$ ) for 10 min. Single-stranded oligonucleotide telomeric probes  $(\text{TTAGGG})_5$  conjugated with Cy3 or biotin were added to hybridization mixture (40% formamide, 2.4 x SSC, and 12% dextran sulphate, 5 ng/ $\mu\text{l}$  labelled probe and 10–50-fold excess of tRNA). Metaphase and lampbrush chromosomes were denatured under a coverslip at 82 C for 5 min. Then slides were incubated with hybridization mixture at room temperature for 12–18 h. After hybridization, slides were washed three times in 2 x SSC at 42°C. Biotin labelled oligonucleotide probes were detected by avidin conjugated with Cy3 (Jackson ImmunoResearch Laboratories). After FISH chromosomal preparations were mounted in DABCO antifade solution containing 1 mg/ml DAPI.

### Immunofluorescent staining of lampbrush chromosomes

Immunostaining of lampbrush chromosome spreads was performed as previously described [32], [33]. For immunostaining we used mouse monoclonal antibodies K121 against 2,2,7-trimethyl guanosine cap (dilution 1:150; Santa Cruz Biotechnology) and rabbit polyclonal antibodies H-300 against coilin (dilution 1:100; Santa Cruz Biotechnology). Lampbrush chromosome spreads were placed in 70%, 50%, 30% ethanol and in PBS with 0.01% Tween-20 for 5 min and blocked in PBS containing 1% blocking reagent (Roche) for 1 h at RT. Slides were incubated with primary antibody for 1 h at RT then washed in PBS, 0.05% Tween-20. The following secondary Abs were used: Cy3-conjugated goat anti-rabbit IgG (dilution 1:500) and Alexa-488-conjugated goat anti-mouse IgG (dilution 1:300; Jackson ImmunoResearch Laboratories). Slides were washed in PBS, 0.05% Tween-20, dehydrated in ethanol series (50%, 70%, 96%) for 5 min, air-dried and mounted in DABCO antifade solution containing 1 mg/ml DAPI.

### Wide-field microscopy

Metaphase and lampbrush chromosomes were examined using Leica fluorescence microscope DM4000 equipped with a monochrome digital camera DFC350 FX and appropriate filter cubes (Leica Wetzlar GmbH, Germany). Images were taken with 10x, 20x, 40x/1 and 100x/1.30 objectives. Leica CW 4000 FISH software was used for acquisition and processing the multicolor images.

### Confocal laser scanning microscopy

For confocal microscopy, nuclei were isolated from oocytes of 0.5–1.5 mm in diameter by jeweler forceps in the isolation medium “5:1” (described above) under the observation at Leica

MZ16 stereomicroscope. Isolated nuclei were incubated for 5 min in “5:1” medium containing 0.07  $\mu$ M Sytox Green (Molecular Probes) [34]. Confocal laser scanning microscopy was carried out with a Leica TCS SP5 microscope based on a Leica DMI 6000 CS inverted microscope. Specimens were examined by the XYZ scanning technique using HC PL APO 20 $\times$  objective and argon laser (496 nm). Images were obtained using LAS AF software (Leica Microsystems, Germany), and 3D reconstruction was processed with Imaris 5.0.1 (Bitplane, AG) software.

## Results


### Genome composition in somatic cells of di- and triploid *P. esculentus* females

In this study we analyzed 27 *P. esculentus* females taken from population systems of R-E type in the Seversky Donets river basin (Eastern Ukraine). The differences between nuclear DNA content of *P. ridibundus* and *P. lessonae* allowed to measure the genome ploidy and to identify the precise genomic composition of hybrid frogs [21], [29] (S1 Table). The individuals with a range of C-values between  $16.00 \pm 35$  were designated as *P. ridibundus*, individuals with a range of C-values between  $14.00 \pm 35$  were designated as *P. lessonae*, while individuals with a range of C-values between  $14.90 \pm 35$  were designated as *P. esculentus* and triploid hybrid frogs with LLR and RRL genotypes have range of C-values between  $21.80 \pm 35$  and  $22.9 \pm 35$  correspondingly [21]. Among analyzed *P. esculentus* females there were 9 diploid (genome composition RL) and 18 triploid (genome composition RRL and LLR) animals. Two frogs had intermediate values of nuclear DNA content (designated as RLX genotype) between corresponding values of nuclear DNA content for LLR and RRL genotypes (S1 Table). Nevertheless unusual types of genomes in oocytes produced by these two frogs allowed us to refer them to RRL genotype (see below).

### Genome composition in oocytes of triploid hybrid frogs with RRL genotype

In R-E population system, we analyzed genome composition in oocytes of 9 diploid, 13 triploid *P. esculentus* females with RRL genotype (including females with RLX genotype) and 5 triploid females with LLR genotype. Algorithm to identify the type of genome transmitted in oocytes of hybrid frogs is represented in S1 Material. At first, we describe the chromosomal sets in oocytes of triploid frogs with RRL genotype. We found that 11 females with RRL genotype produced only one type of oocytes with 13 bivalents corresponding to *P. ridibundus* bivalents (Figs 1a,a' and 2d1–d6'; S1 Movie). Combining the data on hybrid females ploidy and oocyte genome composition, we evaluated appearance of genome elimination and endoreplication events during gametogenesis. We suggest that to form typical oocytes with 13 bivalents corresponding to *P. ridibundus* bivalents, elimination of L genome occurred premeiotically while two remaining R genomes presumably conjugated during meiosis (Fig 3a).

Two females with RLX genotype produced oocytes with unusual chromosomal sets. In one triploid female with RLX genotype (S1 Table), presumably RRL one, 34 oocytes contained 39 univalents, where 26 ones corresponded to *P. ridibundus* lampbrush chromosomes and 13 ones corresponded to *P. lessonae* lampbrush chromosomes (Fig 4b,b'; S1c1–d2 Fig). Importantly, 26 univalents corresponding to *P. ridibundus* lampbrush chromosomes did not form bivalents. Apparently, to form such oocytes neither endoreplication nor elimination occurred during gametogenesis in this triploid female (Fig 5a). In this individual, we also described 4 oocytes with 39 bivalents, where 26 ones were similar to *P. ridibundus* bivalents, while 13 ones were similar to *P. lessonae* bivalents (Fig 4a,a'; S1a1–a6',b1–b3' Fig). Premeiotic endoreplication of the whole




**Fig 1. Typical lampbrush chromosome sets from oocytes of triploid hybrid frogs with RRL and LLR genotypes and diploid hybrid frog with RL genotype.** Full lampbrush chromosome sets from oocytes of triploid hybrid frog with RRL (a,a') and LLR (c,c') genotypes and diploid hybrid frog with RL genotype (b,b'). Chromosome sets are represented by 13 bivalents, which have distribution of marker structures corresponding to *P. ridibundus* (a,a',b,b') or *P. lessonae* (c,c') lampbrush chromosomes. Letter symbols indicate alphabetic numbering of all lampbrush chromosomes; italic type shows correspondence of identified chromosomes to genotype of parental species: *r*—to *P. ridibundus*, *l*—to *P. lessonae*. Chromosomes were counterstained with DAPI (a,b,c). Corresponding phase-contrast micrographs are shown (a',b',c'). Scale bars = 50  $\mu$ m.

doi:10.1371/journal.pone.0123304.g001


triploid karyotype in germ cells without any elimination is required to form oocytes with 39 bivalents (Fig 5a). One oocyte contained 8 bivalents of *P. ridibundus* and 15 univalents corresponding to either *P. ridibundus* or *P. lessonae* lampbrush chromosomes (S2a1–a4 and S3b,b' Figs). Such oocytes indicate abnormalities in conjugation of certain chromosomes of *P. ridibundus* chromosomal set. In that case, individual chromosomes of *P. ridibundus* were lost during oogenesis and endoreplication did not occur (Fig 5a).

For another unusual triploid with RLX genotype, most likely RRL genotype, we obtained 29 oocytes (S1 Table). We found that 13 oocytes contained 13 bivalents, identical to *P. ridibundus*



**Fig 2. Interstitial (TTAGGG)<sub>n</sub> repeat sites mapping allows to identify parental chromosomes in oocytes of hybrid frogs.** (a-c) FISH mapping of (TTAGGG)<sub>n</sub> repeat on metaphase chromosomes of *P. lessonae* (a, a'), *P. ridibundus* (b), and diploid *P. esculentus* (c). One or two interstitial (TTAGGG)<sub>n</sub> repeat sites distinguish parental NOR-bearing chromosomes H (arrows). Asterisks indicate enlarged fragment with two NOR-bearing chromosomes of *P. lessonae*. Arrows indicate interstitial (TTAGGG)<sub>n</sub> repeat sites. (d1–f1') Lampbrush chromosomes from oocytes of triploid hybrid frogs with RRL (d1–d6') and LLR (e1–f1') genotypes. FISH mapping of (TTAGGG)<sub>n</sub> repeat revealed lampbrush chromosome H corresponding to *P. ridibundus* (d6) or *P. lessonae* (e1) LBC H. Interstitial (TTAGGG)<sub>n</sub> repeat sites are shown by square brackets. Lampbrush chromosomes corresponding to *P. ridibundus* LBC F (d1, d1'), G (d2, d2'), D (d3, d3'), I (d4, d4'), B (d5, d5'), and *P. lessonae* LBC B (e2, b2'), F (e3, b3'), L (f1, f1') are shown. Chromosomes on micrographs (d1–d6') were taken from the full lampbrush chromosome set represented on Fig 1a.a'. Chromosomes on micrographs (e1–e3') were taken from the from the full lampbrush chromosome set represented on Fig 1c.c'. Various marker structures are shown by arrows. Chromosomes were counterstained with DAPI. Corresponding phase-contrast micrographs are shown (d1', d2', d3', d4', d5', d6', e1', e2', e3', f1'). Arrowheads indicate centromeres. Scale bars = 10 μm for all panels except a', where scale bar = 2 μm.

doi:10.1371/journal.pone.0123304.g002



**Fig 3. Suggested mechanisms of oogenesis typical for triploid hybrid frogs with RRL genotype and diploid hybrid frogs.** (a) During oogenesis of the majority of triploid hybrids with RRL genotype from studied population systems of R-E type, L genome (blue) was eliminated while two remaining R genomes (brown and orange) without endoreduplication formed 13 bivalents. (b) In oogenesis of the majority of diploid hybrids with RL genotype from studied population systems of R-E type, L genome (blue) was eliminated and the remaining R genome (orange) was endoreduplicated to form 13 bivalents.

doi:10.1371/journal.pone.0123304.g003

bivalents (Fig 5b). Chromosomal sets of the other 15 oocytes from the same frog were represented by 13 bivalents corresponding to *P. ridibundus* lampbrush chromosomes and 13 univalents corresponding to *P. lessonae* lampbrush chromosomes (S3a,a' and S4b1-b4' Figs). Presumably, neither elimination nor endoreduplication were required to form such oocytes during gametogenesis of triploid hybrid female with RRL genotype (Fig 5b). One oocyte from this frog contained 39 bivalents, where segregation of half-bivalents was most likely incomplete.


### Genome composition in oocytes of triploid hybrid frogs with LLR genotype

We determined the genome composition in oocytes from 5 triploid hybrid frogs with LLR genotype (S1 Table). All observed oocytes from 3 triploid hybrid frogs with LLR genotype contained 13 bivalents corresponding to *P. lessonae* bivalents (Figs 1c,c', and 2e1-fl'). To produce oocytes with 13 bivalents corresponding to *P. lessonae* karyotype, R genome had to be eliminated premeiotically while L genome had to form bivalents (S5a Fig).

From oocytes of another triploid hybrid female with LLR genotype we obtained 18 full lampbrush chromosome sets with 13 bivalents corresponding to *P. ridibundus* bivalents (S6b, b' Fig). We suppose that in this hybrid frog, two L genomes were eliminated while the remaining R genome was endoreduplicated premeiotically (S5b Fig).


The majority of chromosomal sets (11 of 16 sets) from the last triploid hybrid female with LLR genotype contained 26 bivalents where 13 ones were similar to *P. ridibundus* chromosomes and 13 ones were similar to *P. lessonae* chromosomes (S3d,d' Fig). Thus, during formation of these oocytes in the triploid frog, one copy of L genome must have been eliminated premeiotically, while remaining L and R genomes must have been endoreduplicated premeiotically (S5c Fig). We also observed 5 lampbrush chromosome sets with 26 univalents, where 13 univalents were similar to *P. ridibundus* chromosomes and 13 univalents were similar to





**Fig 4. Unusual lampbrush chromosome sets from oocytes of triploid hybrid frogs with RRL genotype and two diploid hybrid frogs.** (a,a',b,b') Lampbrush chromosome sets from oocytes of some triploid hybrid frogs with RRL genotype represented by 39 bivalents (a,a') and 39 univalents (b,b'), with 26 bi- or univalents corresponding to *P. ridibundus* lampbrush chromosomes and 13 bi- or univalents corresponding to *P. lessonae* lampbrush chromosomes. (c,c') Lampbrush chromosome set from oocyte of one diploid hybrid frog represented by 26 univalents. Some univalents are similar to *P. ridibundus* lampbrush chromosomes, while other univalents are similar to *P. lessonae* lampbrush chromosomes. (d,d') Lampbrush chromosome set from oocyte of one diploid hybrid frog represented by 26 bivalents corresponding to *P. ridibundus* lampbrush chromosomes. Letter symbols indicate alphabetic numbering of all lampbrush chromosomes; italic type shows correspondence of identified chromosomes to genotype of parental species: *r*—to *P. ridibundus*, *l*—to *P. lessonae*. Chromosomes were counterstained with DAPI (a,b,c,d). Corresponding phase-contrast micrographs are shown (a',b',c',d'). Scale bars = 50  $\mu$ m.

doi:10.1371/journal.pone.0123304.g004



**Fig 5. Suggested additional mechanisms of oogenesis in two triploid frogs with RRL genotype and one diploid hybrid frog.** (a) During oogenesis of one triploid frog with RRL genotype neither elimination nor endoreduplication occurred to form oocytes with 39 univalents (at the top), endoreduplication of all genomes took place to form oocytes with 39 bivalents (in the middle), individual chromosomes from L genome (blue) were lost to form oocytes with aneuploid chromosomal sets (at the bottom). (b) During oogenesis of another triploid frog with RRL genotype elimination of L genome (blue) occurred to form oocytes with 13 bivalents (at the top), premeiotic elimination and endoreduplication were absent to form oocytes with 39 univalents (in the middle), endoreduplication of all genomes took place to form oocytes with 39 bivalents (at the bottom). (c) During oogenesis of one diploid hybrid frog L genome (blue) was eliminated in all observed oocytes. One round of R genome (orange) endoreduplication occurred but bivalents formation was incomplete to form oocytes with both univalents and bivalents (at the top). Two rounds of endoreduplication of R genome took place to form oocytes 26 bivalents (in the middle). One round of R genome endoreduplication occurred but bivalents could not form that led to formation of oocytes with 26 univalents (at the bottom).

doi:10.1371/journal.pone.0123304.g005

*P. lessonae* chromosomes (S3c,c' Fig). To form oocytes with 26 univalents only elimination occurred in germ cells of triploid frog with LLR genotype (S5c Fig).

We conclude that the majority of triploid *P. esculentus* females with RRL and LLR genotypes produced oocytes with 13 bivalents formed by homologous chromosomes, which are represented in double copies in genomes of triploid hybrids. Deviations in genome elimination and existence of additional endoreplication event during oogenesis in triploid hybrid frogs led to formation of oocytes with 26 and 39 bi- or univalents.

## Genome composition in oocytes of diploid hybrid frogs

All oocytes obtained from 5 typical diploid hybrid females contained 13 bivalents corresponding to *P. ridibundus* lampbrush chromosomes (Fig 1b,b'). We suggest that to form oocytes with 13 bivalents, L genome was eliminated while the remaining R genome was endoreplicated premeiotically in diploid hybrid frogs (Fig 3b).

Four diploid *P. esculentus* produced oocytes with different genome composition. In one diploid hybrid female, 20 of the 23 oocytes examined contained 26 bivalents with 13 ones corresponding to *P. lessonae* lampbrush karyotype and 13 ones corresponding to *P. ridibundus* lampbrush karyotype (S4a1-a2' and S6c,c' Figs). To form the oocytes with 26 bivalents in diploid hybrid frog during gametogenesis endoreplication of both L and R genomes was to occur (S7a Fig). Other 3 oocytes from the same frog contained 26 univalents with 13 ones corresponding to *P. lessonae* lampbrush chromosomes and 13 ones corresponding to *P. ridibundus* lampbrush chromosomes (Fig 4c,c'; S4c1-c2' Fig; S2 Movie). To form oocytes with 26 univalents in analyzed diploid hybrid frog, neither elimination nor endoreplication occurred in germ cells (S7a Fig). Previously oocytes with aneuploidy and 26 univalents corresponding to genomes of both parental species were reported for single diploid *P. esculentus* [19].

For another diploid frog, we obtained 6 full lampbrush chromosomal sets represented by 13 bivalents corresponding to *P. ridibundus* lampbrush karyotype. One oocyte from the same frog contained 26 univalents, where 13 univalents were similar to *P. ridibundus* lampbrush chromosomes and other 13 univalents were similar to *P. lessonae* lampbrush chromosomes (S7b Fig).

Among 40 oocytes with full chromosomal sets obtained from another diploid hybrid female, 28 oocytes contained 26 univalents with 13 ones corresponding to *P. ridibundus* chromosomes and 13 ones corresponding to *P. lessonae* chromosomes (S7c Fig). Other 9 oocytes contained various numbers of univalents (15 to 20) which corresponded to lampbrush chromosomes of both parental species. We suppose that aneuploid oocytes may originate after partial loss of chromosomes during gametogenesis without any endoreplication (S7c Fig). Two other sets of lampbrush chromosomes contained 26 bivalents where 13 ones were similar to *P. ridibundus* chromosomes, and 13 ones were similar to *P. lessonae* chromosomes.

In the ovary of the last atypical diploid *P. esculentus* we observed 8 oocytes with 26 bivalents (Fig 4d,d'; S1e1-e2' Fig), 6 oocytes with 26 univalents (S4d1-d2' and S6a,a' Figs) and 16 oocytes with various number of bivalents (from 3 to 10) and univalents (from 8 to 20). Detailed analysis revealed that all examined oocytes contained lampbrush chromosomes corresponding only to *P. ridibundus* karyotype. We suppose that L genome was premeiotically eliminated while R genome was premeiotically endoreplicated ones to form oocytes with 26 univalents and oocytes with both uni- and bivalents and even twice to form oocytes with 26 bivalents (Fig 5c). Alternative premeiotic endoreplication of both R and L genomes and subsequent elimination of doubled L genomes seems to be hardly possible for formation of oocytes with 13 bivalents corresponding to *P. ridibundus* chromosomes. In addition, we have found neither oocytes with lampbrush chromosomes corresponding to *P. lessonae* chromosomes nor aneuploid oocytes which are supposed to occur in such complicated way of oocytes formation. Despite

oocytes with genomes of both parental species were found earlier in diploid *P. esculentus*, oocytes with four identical copies of one parental species genome (26 bivalents of *P. ridibundus*) represent unique data not only for frogs but also for other clonal animals [4], [5], [7], [15], [35], [36]. In oocytes with 26 univalents, identical chromosomes after endoreplication failed to form bivalents. Oocytes with both univalents and bivalents presumably represent unsuccessful attempts to bivalent formation (Fig 5c; S4d1–d2` and S6a,a` Figs).

Therefore, the majority of diploid *P. esculentus* females from Eastern Ukraine population systems of R-E type produced oocytes with 13 bivalents corresponding to *P. ridibundus* chromosomes. We also described diploid hybrid females with deviations in the processes of genome elimination and/or endoreplication, which led to the formation of oocytes with 26 bi- or univalents where genomes of both parental species or only genome of *P. ridibundus* were present.

## Discussion

### Mechanisms of oogenesis in diploid and triploid hybrid European water frogs

Transition to polyploid hybrids creates additional difficulties in gametogenesis, which require changes in the mechanisms of genome elimination and endoreplication. We are the first who represent cytogenetic observations of chromosomal sets from oocytes of triploid European water frogs. These observations are in accordance with the assumption that triploid *P. esculentus* females with RRL and LLR genotypes eliminate premeiotically single copied genome and produce oocytes with remaining genomes [6], [20], [35], [36] (Fig 3a; S5a Fig). 13 bivalents found in oocytes of triploid hybrids are formed between homologous chromosomes of genomes represented in two copies. Thus, our findings confirm the hypothesis suggested by Günther and co-authors in 1979 [20].

Additionally, we established that rare triploid *P. esculentus* females produce variable oocytes with 13 bivalents of single copied genome, oocytes with 26 uni- or bivalents, oocytes with 39 uni- or bivalents and oocytes with both bivalents and univalents (Fig 4a-b`; S3a,a`,c-d` and S6b-b` Figs). Such oocytes allowed us to evaluate occurrence of genome elimination and endoreplication during oogenesis of triploid European water frogs (Fig 5a,b; S5b,c Fig). Genome endoreplication during gametogenesis in triploid females which lead to formation of oocytes with 39 bivalents was not earlier reported for triploid frogs from other population systems. Such abnormal oocytes are common for all parthenogenetic and gynogenetic triploid vertebrate hybrids, which produce unreduced gametes and can develop without fertilization [8], [9], [37]. On the contrary, fertilization is required in the reproduction mode typical for water frogs (hybridogenesis) [16], [17], but the detailed examination of the fate of such oocytes is required. Discussion of the female gametes contribution to the maintenance of the examined European water frog population systems is represented in S2 Material.

It was expected that similar to Central European R-E population systems, in R-E population systems from Eastern Ukraine, R genome is eliminated premeiotically in diploid hybrid frogs. However, in the studied population systems of R-E type, the majority of diploid hybrid females produced oocytes with 13 bivalents of *P. ridibundus* (Fig 3b). We also found no support for previously suggested elimination of R genome or endoreplication of L genome during oogenesis of studied diploid hybrids. The formation of oocytes with 13 bivalents corresponding to *P. ridibundus* chromosomes was discovered for diploid hybrid frogs from Poland population systems of R-L-E type [19]. In contrary to triploids, in diploid hybrids bivalents consist of identical copies appeared after endoreplication event so that recombination can not increase variability in gametes of these animals.

Various deviations from obligatory elimination and endoreplication led to the formation of oocytes with 26 bi- and univalents where genomes of both parental species or only the genome of *P. ridibundus* appeared (Fig 5c; S7a,b,c Fig). Oocytes with 26 bivalents clearly indicate premeiotic endoreplication in diploid *P. esculentus* hybrids. For discussion of diploid frogs role in the maintenance of population system of R-E type see S2 Material.

We also found that the majority of di- and triploid hybrid frogs produced oocytes with variable chromosomal sets (Fig 5a,b,c; S5c and S7a,b,c Figs). We suggest that the processes of genome elimination and endoreplication may occur independently during the development of different germ cells populations. In contrast, previously described diploid males of *P. esculentus* that produced both L and R haploid gametes were considered mosaic [38].

Directed genome duplication in gametogenesis widely spreads among the majority of clonal animals and normally does not happen in sexually reproduced species. In the analysed samples, genome endoreplication occurred premeiotically in few triploid and all diploid hybrid frogs and in one diploid hybrid frog even twice (Figs 3b and 5a,b,c; S5b,c and S7a,b,c Figs). In comparison to elimination event, genome duplication is unselective for any parental species genomes. According to the schemes of the suggested mechanisms of oogenesis (Figs 3b and 5c; S5b,c and S7b Figs) genome endoreplication always occurred after elimination if both processes happened during oogenesis. The omission of cell division after DNA synthesis stage of cell cycle called endoreplication is considered a possible mechanism of genome duplication [9], [39].

## Mechanisms of genome elimination during oogenesis in hybrid European water frogs with different ploidy

Selective genome elimination is a key mechanism leading to appearance of oocytes with one of the parental genomes in water frogs. Possible mechanisms of genome elimination in germ cells of hybrid organisms were previously suggested. It was found that chromosomes of one species may lag and vanish during single [13] or contiguous mitotic [18], [40] or single meiotic division [41]. Differences between sequences of parental species centromeres or centromere binding proteins may be responsible for chromosome loss during division [42]. However, our previous data indicate no any difference in centromere repeats in parental water frog species of geographic origin close to the studied populations [32]. Nevertheless, elimination may also occur during interphase via chromatin budding from the nucleus of germ cells and its further degradation [40], [43].

Not only centromeric but also other repetitive sequences such as transposons differ between closely related parental species [42], [44]. Divergence in transposons in parental genomes leads to the dissimilarities in noncoding RNA in germ cells of hybrid animals playing a role in heterochromatin formation [42]. The mechanism of hybrid disgenesis in *D. melanogaster* mediated by piRNAs selectively blocking activity of transposons from one of the parents [45] may be the mechanism of selective genome elimination in water frog hybrids. We suppose that the genome which was absent and had not transcribed noncoding RNA in maternal oocyte should be eliminated during the gametogenesis in hybrid frog arising after fertilization of this oocyte.

Selective elimination in hybrid frogs may also be similar to paternal genome elimination naturally occurring in some insects and hybridogenetic all-female fishes [46]. In these cases selective genome elimination is supposed to be based on competition between genomes of both parental species for preferential transmitting into gametes. Such genome competition could also lead to elimination-cause mutation in one genome leading to selective elimination of the other parental species genome [46]. Competition between *P. ridibundus* and *P. lessonae* genomes may also result in appearance of elimination-cause mutation more frequently in *P. ridibundus* than in *P. lessonae* genomes.

Oocyte formation in triploid *P. esculentus* frogs is more stable and typically includes only premeiotic genome elimination without endoreplication. Instability in chromatin elimination and appearance of endoreplication lead to formation of oocytes with genome composition being identical to twofold genome composition in somatic cells. Diploid hybrid *P. esculentus* females have more frequent deviations in genome elimination and endoreplication events and can produce oocytes with different genome composition and different ploidy. The majority of oocytes can participate in gamete formation and provide gamete variations required for successful reproduction of di- and triploid hybrid water frogs in different populations.

## Supporting Information

**S1 Fig. Individual chromosome identification in unusual lampbrush chromosome sets.** Identification of individual lampbrush chromosomes from chromosome sets with 39 bivalents (a1–b3') and 39 univalents (c1–d2') from triploid frog with RRL genotype and sets with 26 bivalents (e1–e2') from diploid hybrid frog. (a1–a3) Lampbrush chromosomes corresponding to *P. ridibundus* (a1–a2) or to *P. lessonae* (a3) lampbrush chromosome G. (a4–a6, b1–b3, c1, d1–d2, e1–e2) Lampbrush chromosomes corresponding to *P. ridibundus* (a4–a5', b1–b2', c1–d1', e1–e2') or to *P. lessonae* (a6, a6', b3, b3', d2, d2') lampbrush chromosome H. FISH mapping of (TTAGGG)<sub>n</sub> repeat. Interstitial (TTAGGG)<sub>n</sub> repeat sites are shown by square brackets. Chromosomes on micrographs (a1–a6') were taken from the full chromosome set represented on the [Fig 4a,a'](#). Chromosomes on micrograph (b1–b3') were taken from the other chromosome set with 39 bivalents. Lampbrush chromosomes on micrographs (c1, c1') and (d1–d2') were taken from different chromosome sets containing 39 univalents (full chromosome set not shown and represented on [Fig 4b,b'](#) correspondingly). Chromosomes on micrographs (e1–e2') were taken from the full chromosome set represented on [Fig 4d,d'](#). Various marker structures are shown by arrows. Arrowheads indicate centromeres. Chromosomes were counterstained with DAPI. Corresponding phase-contrast micrographs are shown (a', b', c'). Scale bars = 10 μm. (PDF)

**S2 Fig. Individual chromosome identification in aneuploid lampbrush chromosome set obtained from oocytes of triploid hybrid frog with RRL genotype.** (a1, a2) Bivalent corresponding to *P. ridibundus* lampbrush chromosome H (a1) and univalent corresponding to *P. lessonae* lampbrush chromosome H (a2). (a3, a4) Univalents corresponding to *P. ridibundus* (a3) and *P. lessonae* (a4) lampbrush chromosome B. All chromosomes were taken from the full lampbrush chromosome set represented in [S3b,b'](#) Fig. FISH mapping of (TTAGGG)<sub>n</sub> repeat (a1–a4). Interstitial (TTAGGG)<sub>n</sub> repeat sites are shown by square brackets. Chromosomes were counterstained with DAPI. Arrowheads show centromeres. Scale bars = 10 μm. (PDF)

**S3 Fig. Sample lampbrush chromosome sets from oocytes produced by triploid hybrid females with RRL and LLR genotypes.** (a, a') Lampbrush chromosome set from oocyte of one triploid hybrid female with RRL genotype is represented by 13 bivalents corresponding to *P. ridibundus* chromosomes and 13 univalents corresponding to *P. lessonae* chromosomes. (b, b') Aneuploid lampbrush chromosome set from oocyte of another triploid hybrid female with RRL genotype is represented by 9 bivalents similar to *P. ridibundus* lampbrush chromosomes and about 18 univalents, some of them being similar to *P. ridibundus* lampbrush chromosomes. (c, c') Lampbrush chromosome set from oocyte of one triploid hybrid female with LLR genotype represented by 26 univalents, where 13 univalents correspond to *P. ridibundus* chromosomes and other 13 univalents correspond to *P. lessonae* chromosomes. (d, d') Lampbrush chromosome set from oocyte of another triploid hybrid female with LLR genotype is

represented by 26 bivalents. 13 bivalents are similar to *P. ridibundus* lampbrush chromosomes and 13 bivalents are similar to *P. lessonae* lampbrush chromosomes. Letter symbols indicate alphabetic numbering of all lampbrush chromosomes; italic type shows correspondence of identified chromosomes to genotype of parental species: *r*—to *P. ridibundus*, *l*—to *P. lessonae*. Chromosomes were counterstained with DAPI. Corresponding phase-contrast micrographs are shown (a',b',c',d'). Scale bars = 50  $\mu$ m.

(PDF)

**S4 Fig. Individual chromosome identification in lampbrush chromosome sets with univalents and bivalents.** Lampbrush chromosomes from chromosome sets with 13 bivalents and 13 univalents (b1–b4') from oocytes of triploid hybrid with RRL genotype, 26 bivalents (a1–a2'), and 26 univalents (c1–c2' and d1–d2') from oocytes of different diploid hybrid. Lampbrush chromosomes corresponding to *P. ridibundus* (a1,a1') and to *P. lessonae* (a2,a2') lampbrush chromosome H were taken from full chromosome set represented on [S7c,c' Fig.](#) Bivalents G (b1,b1') and H (b3,b3') are similar to *P. ridibundus* lampbrush chromosomes, and univalents G (b2,b2') and H (b4,b4') are similar to *P. lessonae* lampbrush chromosomes. These lampbrush chromosomes were taken from full lampbrush chromosome set represented on [S3a,a' Fig.](#) Lampbrush chromosomes corresponding to *P. ridibundus* (c1,c1') and to *P. lessonae* (c2,c2') lampbrush chromosome H were taken from full lampbrush chromosome set represented on [Fig 4c,c'](#). Univalents corresponding to *P. ridibundus* lampbrush chromosome H (d1, d1', d2, d2') were taken from chromosome set represented on [S6a,a' Fig.](#) FISH mapping of (TTAGGG)<sub>n</sub> repeat (a1,a2,b3,b4,c1,c2,d1,d2). Interstitial (TTAGGG)<sub>n</sub> repeat sites are shown by square brackets. Chromosomes were counterstained with DAPI. Corresponding phase-contrast micrographs are shown (a1',a2',b3',b4',c1',c2',d1',d2'). Arrows indicate the marker loops. Arrowheads show centromeres. Scale bars = 10  $\mu$ m.

(PDF)

**S5 Fig. Suggested mechanisms of oogenesis in triploid hybrid frogs with LLR genotype.** (a) During oogenesis of triploid hybrid frog with LLR genotype R genome (orange) was eliminated and remaining L genomes (light blue and blue) without endoreplication formed 13 bivalents. (b) During oogenesis of other triploid frog with LLR genotype both L genomes (blue, light blue) were eliminated and R genome was endoreplicated to form oocytes with 13 bivalents. (c) During oogenesis of triploid hybrid frog with LLR genotype elimination of one L genome (light blue) and endoreplication of remaining genomes occurred to form oocytes with 26 bivalents (at the top). Elimination of one L genome (light blue) without endoreplication of remaining genomes took place to form oocytes with 26 univalents (at the bottom).

(PDF)

**S6 Fig. Additional lampbrush chromosome sets obtained from oocytes of diploid hybrid frogs and triploid hybrid frogs with LLR genotype.** (a,a') Lampbrush chromosome set from oocyte of diploid hybrid female represented by 26 univalents corresponding to *P. ridibundus* chromosomes. (b,b') Lampbrush chromosome set from oocyte of diploid hybrid female represented by 13 bivalents corresponding to *P. ridibundus* chromosomes. (c,c') Lampbrush chromosome set from oocyte of triploid hybrid female with LLR genotype represented by 26 bivalents, where 13 bivalents correspond to *P. ridibundus* chromosomes and 13 bivalents correspond to *P. lessonae* chromosomes. Letter symbols indicate alphabetic numbering of all lampbrush chromosomes; italic type shows correspondence of identified chromosomes to genotype of parental species: *r*—to *P. ridibundus*, *l*—to *P. lessonae*. Chromosomes were counterstained with DAPI. Corresponding phase-contrast micrographs are shown (a',b',c'). Scale bars = 50  $\mu$ m.

(PDF)

**S7 Fig. Suggested additional mechanisms of oogenesis in diploid hybrid frogs.** (a) During oogenesis of diploid hybrid frog only endoreplication of both parental genomes occurred to form oocytes with 26 bivalents (at the top), neither elimination nor endoreplication took place to form oocytes with 26 univalents (at the bottom). (b) During oogenesis of other diploid hybrid frog elimination of L genome (blue) and endoreplication of the remaining R genome (orange) occurred to form oocytes with 13 bivalents (at the top), elimination and endoreplication were omitted to form oocytes with 26 univalents (at the bottom). (c) During oogenesis of additional diploid hybrid frog neither elimination nor endoreplication occurred to form oocytes with 26 univalents (at the top). Losing of individual chromosomes corresponding to *P. ridibundus* chromosomes led to formation of aneuploid oocytes (in the middle). Endoreplication took place to form oocytes with 26 bivalents (at the bottom).  
(PDF)

**S8 Fig. Ovaries of parental species and hybrid *P. esculentus* frogs.** Ovary fragments of *P. ridibundus* (a), *P. lessonae* (b), triploid hybrid frog with LLR genotype (c) and diploid hybrid frog (d). Pre-vitellogenic, vitellogenic and post-vitellogenic oocytes (according to Dumont (1972) [\*]) are present in the mature ovaries (a,c) but only pre- and vitellogenic oocytes are present in the immature ovaries (b,d). Ovaries of both parental species (a,b) are characterized by alive oocytes with regular rounded shape and a few dead oocytes. Ovaries of hybrid animals (c,d) have many dead oocytes with irregular shape and abnormal dark brown coloring of oocyte poles. Scale bars = 1 mm. \* Dumont JN (1972) Oogenesis in *Xenopus laevis* (Daudin). I. Stages of oocyte development in laboratory maintained animals. J Morphol 136: 153–180.  
(PDF)

**S9 Fig. Marker structures on lampbrush chromosomes from hybrid frogs.** (a) Detection of chromosome-associated coilin-positive granules by immunofluorescent staining with R288 antibody. (b) Identification of marker loops enriched with splicing factors on lampbrush chromosome corresponding to chromosome I of *P. ridibundus*. Immunofluorescent staining with antibodies against TMG-cap of small nuclear RNA. Arrows show marker loops. Arrowheads indicate centromeres. Chromosomes were counterstained with DAPI. Corresponding phase-contrast micrographs are shown (a',b'). Scale bars = 10 µm.  
(PDF)

**S10 Fig. Chromosome sets of oocytes from females with different genotypes presumably contributing in maintenance of R-E type population systems.** Triploid hybrids with RRL genotype produce oocytes with 13 bivalents corresponding to *P. ridibundus* chromosomes (at the top). Triploid females with LLR genotype produce oocytes with 13 bivalents corresponding to *P. lessonae* chromosomes, oocytes with 13 bivalents corresponding to *P. ridibundus* chromosomes and oocytes with 26 bivalents corresponding to both *P. ridibundus* and *P. lessonae* chromosomes (in the middle). Diploid hybrid frogs produce oocytes with 13 bivalents corresponding to *P. ridibundus* chromosomes, oocytes with 26 bivalents corresponding to both *P. ridibundus* and *P. lessonae* chromosomes and oocytes with 26 bivalents corresponding only to *P. ridibundus* chromosomes (at the bottom).  
(PDF)

**S1 Material. Description of ovaries from parental species and hybrid frogs and algorithm of oocytes karyotype identification.**  
(DOC)



**S2 Material. Mechanisms of hybrid frogs reproduction in the studied population systems of R-E type.**

(DOC)

**S1 Movie. 3D projection of the intact oocyte nucleus from diploid hybrid *P. esculentus* containing lampbrush chromosome set represented by 13 bivalents.**

(AVI)

**S2 Movie. 3D projection of the intact oocyte nucleus from diploid hybrid *P. esculentus* containing lampbrush chromosome set represented by 26 univalents.**

(AVI)

**S1 Table. List of *P. esculentus* females from population systems of R-E type from the Sever-sky Donets river basin in Eastern Ukraine. C values—the amount of DNA per nucleus (genome size, in picograms, pg)—and genotypes are given for each female.**

(PDF)

## Acknowledgments

We are grateful to O. V. Korshunov, O. V. Mykhailova (V. N. Karazin Kharkiv National University, Ukraine) and G. A. Lada (Tambov State University, Russia) for the field assistance. The work was partially performed using experimental equipment of the Research Resource Center 'Chromas' of St Petersburg State University.

## Author Contributions


Conceived and designed the experiments: AK AS. Performed the experiments: DD DS JR SL GM. Analyzed the data: DD SL AK. Contributed reagents/materials/analysis tools: AK SL. Wrote the paper: DD AK. Collected the samples: DD DS.


## References









1. Arnold ML, Hodges SA (1995) Are natural hybrids fit or unfit relative to their parents? *Trends Ecol Evol* 10: 67–71. doi: [10.1016/S0169-5347\(00\)88979-X](https://doi.org/10.1016/S0169-5347(00)88979-X) PMID: [21236955](https://pubmed.ncbi.nlm.nih.gov/21236955/)
2. Chapman MA, Burke JM (2007) Genetic divergence and hybrid speciation. *Evolution* 61: 1773–1780. PMID: [17598755](https://pubmed.ncbi.nlm.nih.gov/17598755/)
3. Kearney M, Fujita MK, Ridenour J (2009) Lost sex in the reptiles: constraints and correlations. In: Schön I, Martens K, van Dijk P, editors. *Lost sex. The evolutionary biology of Parthenogenesis*. Heidelberg, Germany; Springer. pp. 447–474.
4. Lamatsch DK, Stöck M (2009) Sperm-dependent parthenogenesis and hybridogenesis in teleost fishes. In: Schön I, Martens K, van Dijk P, editors. *Lost sex. The evolutionary biology of Parthenogenesis*. Heidelberg, Germany; Springer. pp. 399–432.
5. Bogart JP, Bi K, Fu JZ, Noble DWA, Niedzwiecki J (2007) Unisexual salamanders (genus *Ambystoma*) present a new reproductive mode for eukaryotes. *Genome* 50: 119–136. PMID: [17546077](https://pubmed.ncbi.nlm.nih.gov/17546077/)
6. Plötner J (2005) Die westpaläarktischen Wasserfrösche. Bielefeld: Laurenti-Verlag. 1–161 p. doi: [10.1016/j.ijcard.2010.04.007](https://doi.org/10.1016/j.ijcard.2010.04.007) PMID: [20439124](https://pubmed.ncbi.nlm.nih.gov/20439124/)
7. Stöck M, Ustinova J, Betto-Colliard C, Scharf M, Moritz C, Perrin N (2012) Simultaneous mendelian and clonal genome transmission in a sexually reproducing, all-triploid vertebrate. *Proc R Soc B* 279: 1293–1299. doi: [10.1098/rspb.2011.1738](https://doi.org/10.1098/rspb.2011.1738) PMID: [21993502](https://pubmed.ncbi.nlm.nih.gov/21993502/)
8. Stenberg P, Saura A (2009) Cytology of asexual animals. In: Schön I, Martens K, van Dijk P, editors. *Lost sex. The evolutionary biology of Parthenogenesis*. Heidelberg, Germany; Springer. pp. 63–74.
9. Neaves WB, Baumann P (2011) Unisexual reproduction among vertebrates. *Trends Genet* 27: 81–88. doi: [10.1016/j.tig.2010.12.002](https://doi.org/10.1016/j.tig.2010.12.002) PMID: [21334090](https://pubmed.ncbi.nlm.nih.gov/21334090/)
10. Macgregor HC, Uzzell TM (1964) Gynogenesis in salamanders related to *Ambystoma jeffersonianum*. *Science* 143: 1043–1045. PMID: [14112709](https://pubmed.ncbi.nlm.nih.gov/14112709/)


11. Itono M, Morishima K, Fujimoto T, Bando E, Yamaha E, Arai K (2006) Premeiotic endomitosis produces diploid eggs in the natural clone loach, *Misgurnus anguillicaudatus* (Teleostei: Cobitidae). *J Exp Zool* 305A: 513–523.
12. Lutes AA, Neaves WB, Baumann DP, Wiegraebe W, Baumann P (2010) Sister chromosome pairing maintains heterozygosity in parthenogenetic lizards. *Nature* 464: 283–286. doi: [10.1038/nature08818](https://doi.org/10.1038/nature08818) PMID: [20173738](https://pubmed.ncbi.nlm.nih.gov/20173738/)
13. Cimino MC (1972b) Egg-production, polyploidization and evolution in a diploid all-female fish of the genus *Poeciliopsis*. *Evolution* 26: 294–306.
14. Tunner HG, Heppich S (1981) Premeiotic genome exclusion during oogenesis in the common edible frog *Rana esculenta*. *Naturwissenschaften* 68: 207–208. PMID: [6974310](https://pubmed.ncbi.nlm.nih.gov/6974310/)
15. Alves MJ, Coelho MM, Collares-Pereira MJ (1998) Diversity in the reproductive modes of females of the *Rutilus alburnoides* complex (Teleostei, Cyprinidae): a way to avoid the genetic constraints of uni-parentalism. *Mol Biol Evol* 15: 1233–1242.
16. Berger L (1970) Some characteristics of the crosses within *Rana esculenta* complex in postlarval development. *Ann Zool Fenn* 27: 374–416.
17. Tunner HG (1973) Demonstration of the hybrid origin of the common green frog *Rana esculenta*. *Naturwissenschaften* 60: 481–482. PMID: [4543770](https://pubmed.ncbi.nlm.nih.gov/4543770/)
18. Tunner HG, Heppich-Tunner S (1991) Genome exclusion and two strategies of chromosome duplication in oogenesis of a hybrid frog. *Naturwissenschaften* 78: 32–34.
19. Bucci S, Raghianti M, Mancino G, Berger L, Hotz H, Uzzell T (1990) Lampbrush and mitotic chromosomes of the hemiclonally reproducing hybrid *Rana esculenta* and its parental species. *J Exp Zool* 255: 37–56. PMID: [2391468](https://pubmed.ncbi.nlm.nih.gov/2391468/)
20. Graf JD, Polls-Pelaz M (1989) Evolutionary genetics of the *Rana esculenta* complex. In: Dawley RM, Bogart JP, editors. *Evolution and ecology of unisexual vertebrates*. Albany, New York: New York State Museum Publications. pp. 289–302.
21. Borkin LJ, Korshunov AV, Lada GA, Litvinchuk SN, Rosanov JM, Shabanov DA, et al. (2004) Mass occurrence of polyploid green frogs (*Rana esculenta* complex) in eastern Ukraine. *Russ J Herpetol* 11: 194–213.
22. Shabanov DA, Zinenko AI, Korshunov AV, Kravchenko MO, Mazepa GO (2006) The study of population systems of green frogs (*Rana esculenta* complex) in Kharkov region: history, modern condition and prospects. *Kharkov Nat University J* 3(729): 208–220.
23. Korshunov AV (2009) Ecological patterns of biotopic distribution of *Pelophylax esculentus* complex in Kharkiv region. *Kharkov Nat University J* 8(828): 48–57.
24. Macgregor HC, Sessions SK, Arntzen JW (1990) An integrative analysis of phylogenetic relationships among newts of the genus *Triturus* (family Salamandridae), using comparative biochemistry, cytogenetics and reproductive interactions. *J Evolution Biol* 3: 329–373.
25. Callan HG (1986) *Lampbrush Chromosomes*. London: Springer-Verlag. 1–252 p.
26. Giorgi F, Galleni L (1972) The lampbrush chromosomes of *Rana esculenta* L. (Amphibia-Anura). *Caryologia* 25: 107–123.
27. Uzzell T, Hotz H (1979) Electrophoretic and morphological evidence for two forms of green frogs (*Rana esculenta* complex) in peninsular Italy (Amphibia, Salientia). *Mitt Zool Mus Berlin* 55: 13–27.
28. Graf JD, Müller WP (1979) Experimental gynogenesis provides evidence of hybridogenetic reproduction in the *Rana esculenta* complex. *Experientia* 35: 1574–1576. PMID: [316396](https://pubmed.ncbi.nlm.nih.gov/316396/)
29. Vinogradov AE, Borkin LJ, Günther R, Rosanov JM (1990) Genome elimination in diploid and triploid *Rana esculenta* males: cytological evidence from DNA flow cytometry. *Genome* 33: 619–627. PMID: [2262136](https://pubmed.ncbi.nlm.nih.gov/2262136/)
30. Litvinchuk SN, Borkin LJ, Rosanov JM (2004) Intra- and interspecific genome size variation in hynobiid salamanders of Russia and Kazakhstan: determination by flow cytometry. *Asiat Herpetol Res* 10: 282–294.
31. Gall JG, Murphy C, Callan HG, Wu ZA (1991) Lampbrush chromosomes. *Method Cell Biol* 36: 149–166. PMID: [1811131](https://pubmed.ncbi.nlm.nih.gov/1811131/)
32. Dedukh D, Mazepa G, Shabanov D, Rosanov J, Litvinchuk S, Borkin LJ, et al. (2013) Cytological maps of lampbrush chromosomes of European water frogs (*Pelophylax esculentus* complex) from the Eastern Ukraine. *BMC Genet* 14: 1–26.
33. Krasikova A, Kulikova T, Saifitdinova A, Derjusheva S, Gaginskaya E (2004) Centromeric protein bodies on avian lampbrush chromosomes contain a protein detectable with an antibody against DNA topoisomerase II. *Chromosoma* 113: 316–323. PMID: [15616868](https://pubmed.ncbi.nlm.nih.gov/15616868/)


34. Krasikova A, Khodyuchenko T, Maslova A, Vasilevskaya E (2012) Three-dimensional organisation of RNA-processing machinery in avian growing oocyte nucleus. *Chromosome Research* 20: 979–994. doi: [10.1007/s10577-012-9327-7](https://doi.org/10.1007/s10577-012-9327-7) PMID: [23318709](https://pubmed.ncbi.nlm.nih.gov/23318709/)
35. Pruvost NBM, Hoffmann A, Reyer HU (2013) Gamete production patterns, ploidy, and population genetics reveal evolutionary significant units in hybrid water frogs (*Pelophylax esculentus*). *Ecol Evol* 3(9): 2933–2946. doi: [10.1002/ece3.687](https://doi.org/10.1002/ece3.687) PMID: [24101984](https://pubmed.ncbi.nlm.nih.gov/24101984/)
36. Christiansen DG, Reyer HU (2009) From clonal to sexual hybrids: genetic recombination via triploids in all-hybrid populations of water frogs. *Evolution* 63: 1754–1768. doi: [10.1111/j.1558-5646.2009.00673.x](https://doi.org/10.1111/j.1558-5646.2009.00673.x) PMID: [19245393](https://pubmed.ncbi.nlm.nih.gov/19245393/)
37. Yamashita M, Jiang J, Onozato H, Nakanishi T, Nagahama Y (1993) A tripolar spindle formed at meiosis I assures the retention of the original ploidy in the gynogenetic triploid crucian carp, ginbuna *Carassius auratus langsdorffii*. *Dev Growth Differ* 35: 631–636.
38. Vinogradov AE, Borkin LJ, Günther R, Rosanov JM (1991) Two germ cell lineages with genomes of different species in one and the same animal. *Hereditas* 114(3): 245–251. PMID: [1960101](https://pubmed.ncbi.nlm.nih.gov/1960101/)
39. Edgar BA, Orr-Weaver TL (2001) Endoreplication cell cycles: more for less. *Cell* 105: 297–306. PMID: [11348589](https://pubmed.ncbi.nlm.nih.gov/11348589/)
40. Ogielska M (1994) Nucleus-like bodies in gonial cells of *Rana esculenta* (Amphibia, Anura) tadpoles—a putative way of chromosome elimination. *Zool Pol* 39: 461–474.
41. Zhang Q, Arai K, Yamashita M (1998) Cytogenetic mechanisms for triploid and haploid egg formation in the triploid loach *Misgurnus anguillicaudatus*. *J Exp Zool* 281: 608–619.
42. Brown JD, O'Neill RJ (2010) Chromosomes, conflict, and epigenetics: chromosomal speciation revisited. *Annu Rev Genom Hum G* 11: 291–316. doi: [10.1146/annurev-genom-082509-141554](https://doi.org/10.1146/annurev-genom-082509-141554) PMID: [20438362](https://pubmed.ncbi.nlm.nih.gov/20438362/)
43. Gernand D, Rutten T, Varshney A, Rubtsova M, Prodanovic S, Brüb C, et al. (2005) Uniparental chromosome elimination at mitosis and interphase in wheat and pearl millet crosses involves micronucleus formation, progressive heterochromatinization, and DNA fragmentation. *Plant Cell* 17: 2431–2438. PMID: [16055632](https://pubmed.ncbi.nlm.nih.gov/16055632/)
44. Zale na A, Choleva L, Ogielska M, Rábová M, Marec F, Ráb P (2011) Evidence for integrity of parental genomes in the diploid hybridogenetic water frog *Pelophylax esculentus* by genomic in situ hybridization. *Cytogenet Genome Res* 134: 206–212. doi: [10.1159/000327716](https://doi.org/10.1159/000327716) PMID: [21555873](https://pubmed.ncbi.nlm.nih.gov/21555873/)
45. Brennecke J, Malone CD, Aravin AA, Sachidanandam R, Stark A, Hannon GJ (2008) An epigenetic role for maternally inherited piRNAs in transposon silencing. *Science* 322: 1387–1392. doi: [10.1126/science.1165171](https://doi.org/10.1126/science.1165171) PMID: [19039138](https://pubmed.ncbi.nlm.nih.gov/19039138/)
46. Herrick G, Seger J (1999) Imprinting and paternal genome elimination in insects. In: Ohlsson R, editor. *Genomic imprinting: an interdisciplinary approach*. Heidelberg: Springer. pp. 41–71.








































Female genotype	Oocyte chromosomal set			
RRL	 <p data-bbox="562 644 799 667">13 bivalents of <i>P. ridibundus</i></p>	88,4%		
LLR	 <p data-bbox="562 976 799 999">13 bivalents of <i>P. lessonae</i></p>	60%	 <p data-bbox="1039 960 1294 983">13 bivalents of <i>P. ridibundus</i></p>	 <p data-bbox="1442 954 1883 976">26 bivalents: 13 of <i>P. ridibundus</i>, 13 of <i>P. lessonae</i></p>
RL	 <p data-bbox="562 1272 799 1294">13 bivalents of <i>P. ridibundus</i></p>	75,5%	 <p data-bbox="972 1272 1413 1294">26 bivalents: 13 of <i>P. ridibundus</i>, 13 of <i>P. lessonae</i></p>	 <p data-bbox="1525 1272 1785 1294">26 bivalents of <i>P. ridibundus</i></p>

**Supplementary Table 1.**

**List of *P. esculentus* females from population systems of R-E type from the Seversky Donets river basin**

Frog`s number	Mean of C	Female genotype	Locality	Number of analyzed oocytes	Oocyte chromosomal set	Comment
1	23.14	RRL	Dobritsky prud	1	13 bivalents <i>P.ridibundus</i>	
			Latitude 49°32'	2	13 bivalents <i>P.ridibundus</i>	
			Longitude 36°16'	3	13 bivalents <i>P.ridibundus</i>	
				4	13 bivalents <i>P.ridibundus</i>	
				5	13 bivalents <i>P.ridibundus</i>	
				6	13 bivalents <i>P.ridibundus</i>	
				7	13 bivalents <i>P.ridibundus</i>	
				8	13 bivalents <i>P.ridibundus</i>	
				9	13 bivalents <i>P.ridibundus</i>	
				10	13 bivalents <i>P.ridibundus</i>	
				11	13 bivalents <i>P.ridibundus</i>	
				12	13 bivalents <i>P.ridibundus</i>	
				13	13 bivalents <i>P.ridibundus</i>	
				14	13 bivalents <i>P.ridibundus</i>	
				15	13 bivalents <i>P.ridibundus</i>	
				16	13 bivalents <i>P.ridibundus</i>	
				17	13 bivalents <i>P.ridibundus</i>	
				18	13 bivalents <i>P.ridibundus</i>	
				19	13 bivalents <i>P.ridibundus</i>	
2	23.32	RRL	Dobritsky prud	1	13 bivalents <i>P.ridibundus</i>	
			Latitude 49°32'	2	13 bivalents <i>P.ridibundus</i>	
			Longitude 36°16'	3	13 bivalents <i>P.ridibundus</i>	
				4	13 bivalents <i>P.ridibundus</i>	
				5	13 bivalents <i>P.ridibundus</i>	

				6	13 bivalents <i>P.ridibundus</i>	
				7	13 bivalents <i>P.ridibundus</i>	
				8	13 bivalents <i>P.ridibundus</i>	
				9	13 bivalents <i>P.ridibundus</i>	
				10	13 bivalents <i>P.ridibundus</i>	
				11	13 bivalents <i>P.ridibundus</i>	
				12	13 bivalents <i>P.ridibundus</i>	
				13	13 bivalents <i>P.ridibundus</i>	
				14	13 bivalents <i>P.ridibundus</i>	
3	23.11	RRL	Dobritsky prud Latitude 49°32' Longitude 36°16'	1	13 bivalents <i>P.ridibundus</i>	
				2	13 bivalents <i>P.ridibundus</i>	
				3	13 bivalents <i>P.ridibundus</i>	
				4	13 bivalents <i>P.ridibundus</i>	
				5	13 bivalents <i>P.ridibundus</i>	
				6	13 bivalents <i>P.ridibundus</i>	
				7	13 bivalents <i>P.ridibundus</i>	
				8	13 bivalents <i>P.ridibundus</i>	
				9	13 bivalents <i>P.ridibundus</i>	
				10	13 bivalents <i>P.ridibundus</i>	
				11	13 bivalents <i>P.ridibundus</i>	
				12	13 bivalents <i>P.ridibundus</i>	
				13	13 bivalents <i>P.ridibundus</i>	
				14	13 bivalents <i>P.ridibundus</i>	
				15	13 bivalents <i>P.ridibundus</i>	
				16	13 bivalents <i>P.ridibundus</i>	
				17	13 bivalents <i>P.ridibundus</i>	
				18	13 bivalents <i>P.ridibundus</i>	
				19	13 bivalents <i>P.ridibundus</i>	
				20	13 bivalents <i>P.ridibundus</i>	
				21	13 bivalents <i>P.ridibundus</i>	
				22	13 bivalents <i>P.ridibundus</i>	

4	22.94	RRL	Dobritsky prud Latitude 49°32' Longitude 36°16'	1	13 bivalents <i>P.ridibundus</i>
				2	13 bivalents <i>P.ridibundus</i>
				3	13 bivalents <i>P.ridibundus</i>
				4	13 bivalents <i>P.ridibundus</i>
				5	13 bivalents <i>P.ridibundus</i>
				6	13 bivalents <i>P.ridibundus</i>
				7	13 bivalents <i>P.ridibundus</i>
				8	13 bivalents <i>P.ridibundus</i>
				9	13 bivalents <i>P.ridibundus</i>
				10	13 bivalents <i>P.ridibundus</i>
				11	13 bivalents <i>P.ridibundus</i>
				12	13 bivalents <i>P.ridibundus</i>
				13	13 bivalents <i>P.ridibundus</i>
				14	13 bivalents <i>P.ridibundus</i>
				15	13 bivalents <i>P.ridibundus</i>
				16	13 bivalents <i>P.ridibundus</i>
				17	13 bivalents <i>P.ridibundus</i>
				18	13 bivalents <i>P.ridibundus</i>
				19	13 bivalents <i>P.ridibundus</i>
				20	13 bivalents <i>P.ridibundus</i>
				21	13 bivalents <i>P.ridibundus</i>
				22	13 bivalents <i>P.ridibundus</i>
				23	13 bivalents <i>P.ridibundus</i>
				24	13 bivalents <i>P.ridibundus</i>
				25	13 bivalents <i>P.ridibundus</i>
5	23.25	RRL	Dobritsky prud Latitude 49°32' Longitude 36°16'	1	13 bivalents <i>P.ridibundus</i>
				2	13 bivalents <i>P.ridibundus</i>
				3	13 bivalents <i>P.ridibundus</i>
				4	13 bivalents <i>P.ridibundus</i>
				5	13 bivalents <i>P.ridibundus</i>



				6	13 bivalents <i>P.ridibundus</i>	
				7	13 bivalents <i>P.ridibundus</i>	
				8	13 bivalents <i>P.ridibundus</i>	
				9	13 bivalents <i>P.ridibundus</i>	
				10	13 bivalents <i>P.ridibundus</i>	
				11	13 bivalents <i>P.ridibundus</i>	
				12	13 bivalents <i>P.ridibundus</i>	
				13	13 bivalents <i>P.ridibundus</i>	
				14	13 bivalents <i>P.ridibundus</i>	
				15	13 bivalents <i>P.ridibundus</i>	
				16	13 bivalents <i>P.ridibundus</i>	
				17	13 bivalents <i>P.ridibundus</i>	
				18	13 bivalents <i>P.ridibundus</i>	
6	15,26	RL	Dobritsky prud Latitude 49°32' Longitude 36°16'	1	26 bivalents 13 <i>P.ridibundus</i> , 13 <i>P.lessonae</i>	
				2	26 bivalents 13 <i>P.ridibundus</i> , 13 <i>P.lessonae</i>	
				3	26 bivalents 13 <i>P.ridibundus</i> , 13 <i>P.lessonae</i>	
				4	26 bivalents 13 <i>P.ridibundus</i> , 13 <i>P.lessonae</i>	
				5	26 bivalents 13 <i>P.ridibundus</i> , 13 <i>P.lessonae</i>	
				6	26 bivalents 13 <i>P.ridibundus</i> , 13 <i>P.lessonae</i>	
				7	26 bivalents 13 <i>P.ridibundus</i> , 13 <i>P.lessonae</i>	
				8	26 bivalents 13 <i>P.ridibundus</i> , 13 <i>P.lessonae</i>	
				9	26 univalents 13 <i>P.ridibundus</i> , 13 <i>P.lessonae</i>	
				10	26 univalents 13 <i>P.ridibundus</i> , 13 <i>P.lessonae</i>	
				11	26 bivalents 13 <i>P.ridibundus</i> , 13 <i>P.lessonae</i>	
				12	26 bivalents 13 <i>P.ridibundus</i> , 13 <i>P.lessonae</i>	
				13	26 bivalents 13 <i>P.ridibundus</i> , 13 <i>P.lessonae</i>	
				14	26 bivalents 13 <i>P.ridibundus</i> , 13 <i>P.lessonae</i>	
				15	26 bivalents 13 <i>P.ridibundus</i> , 13 <i>P.lessonae</i>	
				16	26 bivalents 13 <i>P.ridibundus</i> , 13 <i>P.lessonae</i>	FISH with TTAGGG
				17	26 bivalents 13 <i>P.ridibundus</i> , 13 <i>P.lessonae</i>	FISH with TTAGGG
				18	26 bivalents 13 <i>P.ridibundus</i> , 13 <i>P.lessonae</i>	FISH with TTAGGG

				19	26 bivalents 13 P.ridibundus, 13 P.lessonae	
				20	26 bivalents 13 P.ridibundus, 13 P.lessonae	
				21	26 univalents 13 P.ridibundus, 13 P.lessonae	
				22	26 bivalents 13 P.ridibundus, 13 P.lessonae	
				23	26 bivalents 13 P.ridibundus, 13 P.lessonae	
<b>7</b>	22,84	RRL	Pojma (II159) Latitude 49°38' Longitude 36°20'	1	13 bivalents P.ridibundus	
				2	13 bivalents P.ridibundus	
				3	13 bivalents P.ridibundus	
				4	13 bivalents P.ridibundus	
				5	13 bivalents P.ridibundus	
				6	13 bivalents P.ridibundus	
				7	13 bivalents P.ridibundus	
				8	13 bivalents P.ridibundus	
				9	13 bivalents P.ridibundus	
				10	13 bivalents P.ridibundus	
				11	13 bivalents P.ridibundus	
				12	13 bivalents P.ridibundus	
				13	13 bivalents P.ridibundus	
				14	13 bivalents P.ridibundus	
				15	13 bivalents P.ridibundus	
				16	13 bivalents P.ridibundus	
				17	13 bivalents P.ridibundus	
				18	13 bivalents P.ridibundus	
				19	13 bivalents P.ridibundus	
				20	13 bivalents P.ridibundus	
<b>8</b>	23,01	RRL	Pojma (II188) Latitude 49°38' Longitude 36°20'	1	13 bivalents P.ridibundus	
				2	13 bivalents P.ridibundus	
				3	13 bivalents P.ridibundus	
				4	13 bivalents P.ridibundus	
				5	13 bivalents P.ridibundus	

				6	13 bivalents <i>P.ridibundus</i>	
				7	13 bivalents <i>P.ridibundus</i>	
				8	13 bivalents <i>P.ridibundus</i>	
				9	13 bivalents <i>P.ridibundus</i>	
				10	13 bivalents <i>P.ridibundus</i>	
				11	13 bivalents <i>P.ridibundus</i>	
				12	13 bivalents <i>P.ridibundus</i>	
				13	13 bivalents <i>P.ridibundus</i>	
				14	13 bivalents <i>P.ridibundus</i>	
				15	13 bivalents <i>P.ridibundus</i>	
				16	13 bivalents <i>P.ridibundus</i>	
				17	13 bivalents <i>P.ridibundus</i>	
				18	13 bivalents <i>P.ridibundus</i>	
				19	13 bivalents <i>P.ridibundus</i>	
				20	13 bivalents <i>P.ridibundus</i>	
9	22,15	LRx	Pojma (II162) Latitude 49°38' Longitude 36°20'	1	39 univalents	FISH with TTAGGG
				2	39 univalents	
				3	39 bivalents 26 <i>P.ridibundus</i> , 13 <i>P.lessonae</i>	
				4	39 univalents 26 <i>P.ridibundus</i> , 13 <i>P.lessonae</i>	
				5	39 univalents	
				6	39 univalents 26 <i>P.ridibundus</i> , 13 <i>P.lessonae</i>	
				7	39 univalents	
				8	39 univalents	
				9	39 univalents	
				10	39 univalents	
				11	39 univalents	
				12	39 univalents 26 <i>P.ridibundus</i> , 13 <i>P.lessonae</i>	
				13	39 univalents	
				14	39 univalents	
				15	39 univalents	
				16	39 univalents	

				17	39 univalents	
				18	39 univalents	
				19	39 univalents	
				20	aneuploid, bi- and univalents	FISH with TTAGGG
				21	39 univalents	
				22	39 univalents	
				23	39 univalents	
				24	39 univalents	
				25	39 univalents	
				26	39 univalents	
				27	39 univalents 26 P.ridibundus, 13 P.lessonae	
				28	39 univalents	
				29	39 univalents	
				30	39 univalents 26 P.ridibundus, 13 P.lessonae	
				31	39 univalents 26 P.ridibundus, 13 P.lessonae	FISH with TTAGGG
				32	39 univalents 26 P.ridibundus, 13 P.lessonae	FISH with TTAGGG
				33	39 univalents 26 P.ridibundus, 13 P.lessonae	FISH with TTAGGG
				34	39 univalents	
				35	39 univalents	
				36	39 bivalents 26 P.ridibundus, 13 P.lessonae	FISH with TTAGGG
				37	39 bivalents 26 P.ridibundus, 13 P.lessonae	FISH with TTAGGG
				38	39 bivalents 26 P.ridibundus, 13 P.lessonae	FISH with TTAGGG
				39	39 univalents	FISH with TTAGGG
10	22,35	LRx	Pojma (II165) Latitude 49°38' Longitude 36°20'	1	13 bivalents P.ridibundus, 13 univalents P.lessonae	FISH with TTAGGG
				2	13 bivalents P.ridibundus	
				3	13 bivalents P.ridibundus	
				4	39 bivalents 26 P.ridibundus, 13 P.lessonae	FISH with TTAGGG
				5	13 bivalents P.ridibundus	FISH with TTAGGG
				6	13 bivalents P.ridibundus, 13 univalents P.lessonae	FISH with TTAGGG
				7	13 bivalents P.ridibundus, 13 univalents P.lessonae	FISH with TTAGGG
				8	13 bivalents P.ridibundus, 13 univalents P.lessonae	FISH with TTAGGG

				9	13 bivalents <i>P.ridibundus</i> , 13 univalents <i>P.lessonae</i>	FISH with TTAGGG
				10	13 bivalents <i>P.ridibundus</i> , 13 univalents <i>P.lessonae</i>	FISH with TTAGGG
				11	13 bivalents <i>P.ridibundus</i>	FISH with TTAGGG
				12	13 bivalents <i>P.ridibundus</i>	
				13	13 bivalents <i>P.ridibundus</i> , 13 univalents <i>P.lessonae</i>	
				14	13 bivalents <i>P.ridibundus</i> , 13 univalents <i>P.lessonae</i>	
				15	13 bivalents <i>P.ridibundus</i>	
				16	13 bivalents <i>P.ridibundus</i> , 13 univalents <i>P.lessonae</i>	
				17	13 bivalents <i>P.ridibundus</i>	
				18	13 bivalents <i>P.ridibundus</i> , 13 univalents <i>P.lessonae</i>	
				19	13 bivalents <i>P.ridibundus</i> , 13 univalents <i>P.lessonae</i>	
				20	13 bivalents <i>P.ridibundus</i> , 13 univalents <i>P.lessonae</i>	
				21	13 bivalents <i>P.ridibundus</i> , 13 univalents <i>P.lessonae</i>	
				22	13 bivalents <i>P.ridibundus</i> , 13 univalents <i>P.lessonae</i>	
				23	13 bivalents <i>P.ridibundus</i>	
				24	13 bivalents <i>P.ridibundus</i>	
				25	13 bivalents <i>P.ridibundus</i>	
				26	13 bivalents <i>P.ridibundus</i>	
				27	13 bivalents <i>P.ridibundus</i> , 13 univalents <i>P.lessonae</i>	
				28	13 bivalents <i>P.ridibundus</i>	
				29	13 bivalents <i>P.ridibundus</i>	
<b>11</b>	15,17	RL	Pojma (II168) Latitude 49°38' Longitude 36°20'	1	26 bivalents <i>P.ridibundus</i>	FISH with TTAGGG
				2	26 univalents <i>P.ridibundus</i>	FISH with TTAGGG
				3	26 bivalents <i>P.ridibundus</i>	FISH with TTAGGG
				4	26 univalents <i>P.ridibundus</i>	FISH with TTAGGG
				5	26 univalents <i>P.ridibundus</i>	FISH with TTAGGG
				6	26 bivalents <i>P.ridibundus</i>	FISH with TTAGGG
				7	26 bivalents <i>P.ridibundus</i>	FISH with TTAGGG
				8	26 univalents <i>P.ridibundus</i>	FISH with TTAGGG
				9	26 bivalents <i>P.ridibundus</i>	FISH with TTAGGG
				10	4 bivalents, 15 univalents <i>P.ridibundus</i>	FISH with TTAGGG

				11	6 bivalents, 14 univalents P.ridibundus	FISH with TTAGGG
				12	3 bivalents, 19 univalents P.ridibundus	FISH with TTAGGG
				13	3 bivalents, 16 univalents P.ridibundus	
				14	4 bivalents, 15 univalents P.ridibundus	
				15	10 bivalents, 6 unival, P.ridibundus	
				16	26 bivalents P.ridibundus	
				17	26 bivalents P.ridibundus	
				18	4 bivalents, 14 univalents P.ridibundus	
				19	3 bivalents, 15 univalents P.ridibundus	
				20	5 bivalents, 10 univalents P.ridibundus	
				21	3 bivalents, 19 univalents P.ridibundus	
				22	5 bivalents, 16 univalents P.ridibundus	
				23	5 bivalents, 16 univalents P.ridibundus	
				24	5 bivalents, 15 univalents P.ridibundus	
				25	3 bivalents, 29 univalents P.ridibundus	
				26	26 bivalents P.ridibundus	
				27	26 bivalents P.ridibundus	
				28	26 univalents P.ridibundus	
				29	4 bivalents, 18 univalents P.ridibundus	
				30	26 univalents P.ridibundus	
				31	3 bivalents, 19 univalents P.ridibundus	
12	20,28	RLL	Iskov prud Latitude 49°33' Longitude 36°17'	1	26 univalents 13 P.ridibundus, 13 P.lessonae	
				2	26 univalents 13 P.ridibundus, 13 P.lessonae	
				3	26 bivalents 13 P.ridibundus, 13 P.lessonae	
				4	26 univalents 13 P.ridibundus, 13 P.lessonae	FISH with TTAGGG
				5	26 bivalents 13 P.ridibundus, 13 P.lessonae	FISH with TTAGGG
				6	26 bivalents 13 P.ridibundus, 13 P.lessonae	FISH with TTAGGG
				7	26 bivalents 13 P.ridibundus, 13 P.lessonae	
				8	26 bivalents 13 P.ridibundus, 13 P.lessonae	
				9	26 univalents 13 P.ridibundus, 13 P.lessonae	
				10	26 bivalents 13 P.ridibundus, 13 P.lessonae	

				11	26 bivalents 13 P.ridibundus, 13 P.lessonae	
				12	26 bivalents 13 P.ridibundus, 13 P.lessonae	
				13	26 bivalents 13 P.ridibundus, 13 P.lessonae	
				14	26 bivalents 13 P.ridibundus, 13 P.lessonae	
				15	26 univalents 13 P.ridibundus, 13 P.lessonae	
				16	26 bivalents 13 P.ridibundus, 13 P.lessonae	
<b>13</b>	14,65	RL	Iskov prud Latitude 49°33' Longitude 36°17'	1	13 bivalents P.ridibundus	
				2	13 bivalents P.ridibundus	
				3	13 bivalents P.ridibundus	
				4	13 bivalents P.ridibundus	
				5	26 univalents 13 P.ridibundus, 13 P.lessonae	
				6	13 bivalents P.ridibundus	
				7	13 bivalents P.ridibundus	
<b>14</b>	14,92	RL	Iskov prud (H5) Latitude 49°33' Longitude 36°17'	1	13 bivalents P.ridibundus	
				2	13 bivalents P.ridibundus	
				3	13 bivalents P.ridibundus	
				4	13 bivalents P.ridibundus	
				5	13 bivalents P.ridibundus	
				6	13 bivalents P.ridibundus	
				7	13 bivalents P.ridibundus	
				8	13 bivalents P.ridibundus	
				9	13 bivalents P.ridibundus	
				10	13 bivalents P.ridibundus	
				11	13 bivalents P.ridibundus	
				12	13 bivalents P.ridibundus	
				13	13 bivalents P.ridibundus	
				14	13 bivalents P.ridibundus	
				15	13 bivalents P.ridibundus	
				16	13 bivalents P.ridibundus	
				17	13 bivalents P.ridibundus	

				18	13 bivalents P.ridibundus	immunostaining against coilin
				19	13 bivalents P.ridibundus	immunostaining against coilin
				20	13 bivalents P.ridibundus	immunostaining against coilin
				21	13 bivalents P.ridibundus	
				22	13 bivalents P.ridibundus	immunostaining with K 121
				23	13 bivalents P.ridibundus	immunostaining with K 121
<b>15</b>	22,88	RRL	Sykhaja Gomolsha (H Latitude 49°32' Longitude 36°20'	1 2 3 4 5	13 bivalents P.ridibundus 13 bivalents P.ridibundus 13 bivalents P.ridibundus 13 bivalents P.ridibundus 13 bivalents P.ridibundus	
<b>16</b>	22,68	RRL	Sykhaja Gomolsha (4 Latitude 49°32' Longitude 36°20'	1 2 3 4 5 6 7 8 9 10 11 12 13 14 15 16	13 bivalents P.ridibundus 13 bivalents P.ridibundus 13 bivalents P.ridibundus 13 bivalents P.ridibundus 13 bivalents P.ridibundus 13 bivalents P.ridibundus 13 bivalents P.ridibundus 13 bivalents P.ridibundus 13 bivalents P.ridibundus 13 bivalents P.ridibundus 13 bivalents P.ridibundus 13 bivalents P.ridibundus 13 bivalents P.ridibundus 13 bivalents P.ridibundus 13 bivalents P.ridibundus 13 bivalents P.ridibundus	
<b>17</b>	22,75	RRL	Sykhaja Gomolsha (H Latitude 49°32'	1 2	13 bivalents P.ridibundus 13 bivalents P.ridibundus	



			Longitude 36°20'	3	13 bivalents P.ridibundus	
				4	13 bivalents P.ridibundus	
				5	13 bivalents P.ridibundus	
				6	13 bivalents P.ridibundus	
				7	13 bivalents P.ridibundus	
				8	13 bivalents P.ridibundus	
				9	13 bivalents P.ridibundus	
				10	13 bivalents P.ridibundus	
				11	13 bivalents P.ridibundus	
				12	13 bivalents P.ridibundus	
				13	13 bivalents P.ridibundus	
				14	13 bivalents P.ridibundus	
				15	13 bivalents P.ridibundus	
				16	13 bivalents P.ridibundus	
				17	13 bivalents P.ridibundus	
				18	13 bivalents P.ridibundus	
				19	13 bivalents P.ridibundus	
				20	13 bivalents P.ridibundus	
				21	13 bivalents P.ridibundus	
				22	13 bivalents P.ridibundus	
				23	13 bivalents P.ridibundus	
				24	13 bivalents P.ridibundus	
<b>18</b>	23,01	RRL	Sykhaja Gomolsha (C Latitude 49°32' Longitude 36°20'	1	13 bivalents P.ridibundus	
				2	13 bivalents P.ridibundus	
				3	13 bivalents P.ridibundus	
				4	13 bivalents P.ridibundus	
				5	13 bivalents P.ridibundus	
				6	13 bivalents P.ridibundus	
				7	13 bivalents P.ridibundus	
				8	13 bivalents P.ridibundus	
				9	13 bivalents P.ridibundus	

				10	13 bivalents P.ridibundus	
<b>19</b>	15,01	RL	Sykhaja Gomolsha (C Latitude 49°32' Longitude 36°20'	1	13 bivalents P.ridibundus	
				2	13 bivalents P.ridibundus	
				3	13 bivalents P.ridibundus	
				4	13 bivalents P.ridibundus	
<b>20</b>	14,91	RL	Sykhaja Gomolsha (2 Latitude 49°32' Longitude 36°20'	1	13 bivalents P.ridibundus	
				2	13 bivalents P.ridibundus	immunostaining against coilin
				3	13 bivalents P.ridibundus	immunostaining against coilin
				4	13 bivalents P.ridibundus	immunostaining against coilin
				5	13 bivalents P.ridibundus	
<b>21</b>	15,02	RL	Sykhaja Gomolsha (E Latitude 49°32' Longitude 36°20'	1	13 bivalents P.ridibundus	
				2	13 bivalents P.ridibundus	
				3	13 bivalents P.ridibundus	
				4	13 bivalents P.ridibundus	
				5	13 bivalents P.ridibundus	
				6	13 bivalents P.ridibundus	
				7	13 bivalents P.ridibundus	
				8	13 bivalents P.ridibundus	
				9	13 bivalents P.ridibundus	
				10	13 bivalents P.ridibundus	
				11	13 bivalents P.ridibundus	
				12	13 bivalents P.ridibundus	
				13	13 bivalents P.ridibundus	
				14	13 bivalents P.ridibundus	
				15	13 bivalents P.ridibundus	
				16	13 bivalents P.ridibundus	
				17	13 bivalents P.ridibundus	



				8	13 bivalents P.ridibundus	
				9	13 bivalents P.ridibundus	
				10	13 bivalents P.ridibundus	
				11	13 bivalents P.ridibundus	
				12	13 bivalents P.ridibundus	
				13	13 bivalents P.ridibundus	
				14	13 bivalents P.ridibundus	
				15	13 bivalents P.ridibundus	
				16	13 bivalents P.ridibundus	
				17	13 bivalents P.ridibundus	
				18	13 bivalents P.ridibundus	
				19	13 bivalents P.ridibundus	
				20	13 bivalents P.ridibundus	
				21	13 bivalents P.ridibundus	
				22	13 bivalents P.ridibundus	
				23	13 bivalents P.ridibundus	
<b>25</b>	21,87	RLL	Krasnoyarskoe Lake Latitude 49°01' Longitude 37°68'	1	13 bivalents P.lessonae	
				2	13 bivalents P.lessonae	
				3	13 bivalents P.lessonae	
				4	13 bivalents P.lessonae	
				5	13 bivalents P.lessonae	
				6	13 bivalents P.lessonae	
				7	13 bivalents P.lessonae	
				8	13 bivalents P.lessonae	
				9	13 bivalents P.lessonae	
<b>26</b>	21,73	RLL	Krasnoyarskoe Lake Latitude 49°01' Longitude 37°68'	1	13 bivalents	
				2	13 bivalents	
				3	13 bivalents	
				4	13 bivalents	
				5	13 bivalents P.lessonae	

				6	13 bivalents P.lessonae	
				7	13 bivalents P.lessonae	
				8	13 bivalents P.lessonae	
				9	13 bivalents P.lessonae	
				10	13 bivalents P.lessonae	
				11	13 bivalents P.lessonae	
27	15,06	RL	Krasnoyarskoe Lake Latitude 49°01' Longitude 37°68'	1	26 univalents 13 P.ridibundus, 13 P.lessonae	
				2	26 univalents 13 P.ridibundus, 13 P.lessonae	
				3	26 univalents 13 P.ridibundus, 13 P.lessonae	
				4	26 univalents 13 P.ridibundus, 13 P.lessonae	
				5	26 univalents 13 P.ridibundus, 13 P.lessonae	
				6	26 univalents 13 P.ridibundus, 13 P.lessonae	
				7	aneuploid, 20 univalents	
				8	26 univalents 13 P.ridibundus, 13 P.lessonae	
				9	26 univalents 13 P.ridibundus, 13 P.lessonae	
				10	26 univalents 13 P.ridibundus, 13 P.lessonae	
				11	26 univalents 13 P.ridibundus, 13 P.lessonae	
				12	aneuploid, 15 univalents	
				13	26 univalents 13 P.ridibundus, 13 P.lessonae	
				14	aneuploid, 20 univalents	
				15	26 univalents 13 P.ridibundus, 13 P.lessonae	
				16	26 univalents 13 P.ridibundus, 13 P.lessonae	
				17	26 univalents 13 P.ridibundus, 13 P.lessonae	
				18	26 univalents 13 P.ridibundus, 13 P.lessonae	FISH with TTAGGG
				19	26 univalents 13 P.ridibundus, 13 P.lessonae	FISH with TTAGGG
				20	26 univalents 13 P.ridibundus, 13 P.lessonae	FISH with TTAGGG
				21	26 univalents 13 P.ridibundus, 13 P.lessonae	immunostaining with K 121
				22	aneuploid, 18 univalents	immunostaining with K 121
				23	26 bivalents 13 P.ridibundus, 13 P.lessonae	immunostaining with K 121
				24	26 univalents 13 P.ridibundus, 13 P.lessonae	immunostaining with K 121
				25	aneuploid, 18 univalents	

			26	26 univalents 13 P.ridibundus, 13 P.lessonae	immunostaining against coilin
			27	26 univalents 13 P.ridibundus, 13 P.lessonae	
			28	26 univalents 13 P.ridibundus, 13 P.lessonae	
			29	26 univalents 13 P.ridibundus, 13 P.lessonae	
			30	26 univalents 13 P.ridibundus, 13 P.lessonae	
			31	26 univalents 13 P.ridibundus, 13 P.lessonae	
			32	aneuploid, 23 univalents	
			33	26 univalents 13 P.ridibundus, 13 P.lessonae	
			34	26 univalents 13 P.ridibundus, 13 P.lessonae	
			35	aneuploid, 20 univalents	
			36	26 bivalents 13 P.ridibundus, 13 P.lessonae	
			37	aneuploid, 20 univalents	
			38	26 univalents 13 P.ridibundus, 13 P.lessonae	
			39	26 univalents 13 P.ridibundus, 13 P.lessonae	
			40	aneuploid, 22 univalents	

### Legend

- very good chromosomal preparations. All chromosomes lay separately and they are clearly identified
- good chromosomal prepararions. Individual chromosomes can be overlaped or stretched, nearly all chromosomes are well identified
- chromosomal preparations of average quality. Some chromosomes are stretched or destroyed, the majority chromosomes are identified

# **S1 Material. Supplementary Results. Description of ovaries from parental species and hybrid frogs and algorithm of lampbrush chromosome identification.**

## **Morphology of ovaries from parental and hybrid species**

To evaluate fertility of hybrid and parental females we analysed ovaries according to Ogielska et al. [1]. In comparison to parental species ovaries of hybrid animals had many large perished oocytes which had irregular shape and lacked differential pole colouring. In some hybrid frogs, proportion of viable oocytes was about 20-30% from all mature oocytes; such oocytes can most likely become gametes (S8 Fig.).

## **Chromosome sets from oocytes of hybrid frogs**

To study mechanisms of gametogenesis in di- and triploid hybrid females we analyzed chromosome sets from growing oocytes. The identification of lampbrush chromosomes (LBCs) obtained from oocytes of hybrids was based on their cytological maps available for both parental species individuals of close geographic origin [2]. In oocytes of hybrid frogs, marker structures on LBCs B, D, E, F, G, H and I were most reliable for morphological identification of *P. ridibundus* chromosomes (Figs. 1a,a` and 2d1-2d6`), while LBCs B, C, E, F, G, H and L were most reliable for morphological identification of *P. lessonae* chromosomes (Figs. 1c,c` and 2e1-2f1`). Other LBCs were unreliable for morphological identification of parental species karyotypes, since marker structures in these chromosomes either were formed in similar positions in both parental species or were highly polymorphic.

To simplify lampbrush chromosomes assignment to karyotype of one of the parental species, we identified landmark structures such as marker loops accumulating splicing factors and coilin-

containing spheres and granules using immunofluorescent staining with antibodies against specific components (S9a,a',b,b' Fig.). To confirm the cytological identification, FISH with probe to telomeric (TTAGGG)<sub>n</sub> repeat was performed on lampbrush chromosome sets from oocytes of nearly all hybrid frogs. Mitotic chromosomes represented in Fig. 2a,b clearly demonstrate dissimilarities in interstitial telomere repeat sites (ITSs) on NOR-bearing chromosomes in parental species. Such differences enable to discriminate corresponding chromosomes in genomes of hybrid animals (Fig. 2c). In lampbrush chromosomes set analysis this approach allowed to identify and to assign accurately the NOR-bearing LBC H, which differs in two parental species in the positions of ITSs [2] (Fig. 2d,e1).

Previously we found that chromosome associated nucleolus forms at the nucleolus organizer region on LBC H in oocytes of *P. ridibundus* but not *P. lessonae* from the Seversky Donets river basin, representing a promising marker for identification of the parental species karyotypes in hybrid frogs [2]. Here we demonstrate that in some hybrids, in oocytes where genomes of both parental species were present (oocytes with 26 and 39 uni- or bivalents), nucleolus appeared on lampbrush chromosome corresponding to *P. lessonae* LBC H (S1a6,a6',b3,b3',d2,d2', S2a2 and S4a2,a2',c2,c2' Figs. ). At the same time, nucleolus that formed on lampbrush chromosome corresponding to *P. ridibundus* LBC H was sometimes small and even undetectable (Fig. 1a4,a4',b1,b1'; S4b3,b3',c1,c1' Fig.). Inactivation of the nucleolus organizer region in such oocytes seems to be accidental, but may be mediated by long or short noncoding RNAs transcribed from *P. lessonae* genome.



## Supplementary references

1. Ogielska M, Rozenblut B, Augustynska R, Kotusz A (2010) Degeneration of germ line cells in amphibian ovary. *Acta Zool Stockholm* 91: 319–327.
2. Dedukh D, Mazepa G, Shabanov D, Rosanov J, Litvinchuk S, Borkin LJ, et al. (2013) Cytological maps of lampbrush chromosomes of European water frogs (*Pelophylax esculentus* complex) from the Eastern Ukraine. *BMC Genet* 14: 1–26.

## **Material S2. Supplementary Discussion. Mechanisms of hybrid frogs reproduction in the studied population systems of R-E type.**

We further evaluated the contribution of female gametes to the maintenance of the examined European water frog population systems of R-E type from Eastern Ukraine and suggested mechanisms of hybrid frogs reproduction (S10 Fig.). In S10 Fig. we represent all identified genomes transmitted in oocyte of diploid and triploid hybrid frogs which are supposed to form fertile gametes.

In *P. esculentus* population system of R-E type, oocytes with 13 bivalents produced by the majority of diploid, triploid RRL and one exceptional LLR hybrids can undergo meiosis and form haploid gametes containing *P. ridibundus* genome (S10 Fig.). Our results are opposite to previous studies of R-E population system from Poland and Germany where hybrid frogs which usually produce L gametes lead to hybrid frogs appearance after crossing with parental species (*P. ridibundus*) [1]–[3]. The majority of studied females with LLR genotypes could produce haploid gametes with *P. lessonae* genome. Such oocytes constitute about 60% among all analyzed oocytes from triploid frogs with LLR genotype (S10 Fig.). In addition, triploid frogs with LLR genotype produced oocytes with 13 bivalents corresponding to *P. ridibundus* chromosomes (20%) and oocytes with 26 bivalents corresponding to both *P. ridibundus* and *P. lessonae* genomes (about 14%).

According to our data the appearance of numerous diploid hybrid frogs in population system of R-E type is possible when haploid eggs with R genome are fertilized by the haploid sperm that contains L genome and was presumably produced by hybrid males, because mature *P. lessonae* individuals were not found in the examined population systems [4]. Fertilization of haploid egg with R genome by haploid sperm with R genome presumably gives rise to *P. ridibundus* parental

species. Thus, in the examined population system of R-E type, genome composition presumably differs between haploid gametes in male and female hybrids.

We suppose that oocytes with 39 bivalents produced by triploid frogs are inessential for the maintenance of *P. esculentus* population systems in Eastern Ukraine since mature tetraploid hybrids were not reported in the population systems of R-E, L-E or E types. Furthermore, oocytes with univalents probably cannot overcome meiosis to produce fertile gametes. Oocytes with univalents and 39 bivalents composed about 12% among all observed oocytes produced by triploid frogs with RRL genotype (S10 Fig.). Several hybrid frogs produced oocytes with 26 bivalents where *P. ridibundus* and *P. lessonae* genomes or only *P. ridibundus* genome were present that constitute about 13% among all analyzed oocytes of diploid hybrid frogs (S10 Fig.). Such oocytes could complete meiosis and form diploid gametes with both *P. ridibundus* and *P. lessonae* genomes or two *P. ridibundus* genomes.

After fertilization by haploid sperm with R or L genomes [5], oocytes with 26 bivalents produced by some diploid *P. esculentus* may give rise to triploid RRL or LLR hybrids. Nevertheless, the number of diploid eggs with R or L genomes is presumably insufficient to sustain such a number of triploid hybrids in the population systems from Eastern Ukraine (up to 80% according to [6]). We conclude that diploid sperm is required to produce a large number of triploid hybrids in the analyzed populations of water frogs.

## Supplementary references

1. Plötner, J (2005) Die westpaläarktischen Wasserfrösche. Bielefeld: Laurenti-Verlag. 1–161 p.
2. Graf JD, Polls-Pelaz M (1989) Evolutionary genetics of the *Rana esculenta* complex. In: Dawley RM, Bogart JP, editors. Evolution and ecology of unisexual vertebrates. Albany, New York: New York State Museum Publications. pp. 289–302.
3. Berger L (1990) On the origin of genetic systems in European water frog hybrids. Zool Pol 35(1–4): 5–32.
4. Korshunov AV (2009) Ecological patterns of biotopic distribution of *Pelophylax esculentus* complex in Kharkiv region. Kharkov Nat University J 8(828): 48–57.
5. Borkin LJ, Zinenko AI, Korshunov AV, Lada GA, Litvinchuk SN, Rosanov JM, et al. (2005) Mass polyploidy in *Rana esculenta* hybridogenic complex (Ranidae, Anura, Amphibia) in eastern Ukraine. Proceedings of the 1th Conference of the Ukrainian Herpetological Society. Kyiv. pp. 23–26.
6. Borkin LJ, Korshunov AV, Lada GA, Litvinchuk SN, Rosanov JM, Shabanov DA, et al. (2004) Mass occurrence of polyploid green frogs (*Rana esculenta* complex) in eastern Ukraine. Russ J Herpetol 11: 194–213.